

1 **Grain harvesting as a local source of *Cladosporium* spp. in Denmark**

2

3 Yulia Olsen*, Tanja Begovic, Carsten Ambelas Skjøth, Karen Rasmussen, Ulrich Gosewinkel, Ole Hertel, Torben
4 Sigsgaard

5

6 Yulia Olsen; Torben Sigsgaard

7 *Department of Public Health, Aarhus University, Bartholins Allé 2, 8000 Aarhus, Denmark*

8

9 Tanja Begovic; Ulrich Gosewinkel

10 *Department of Environmental Science – Environmental microbiology & biotechnology, Aarhus University,*

11 *Frederiksborgvej 399, 4000 Roskilde, Denmark*

12

13 Carsten Ambelas Skjøth

14 *School of Science and the Environment, University of Worcester, Henwick Grove, Worcester WR2 6AJ, UK*

15

16 Karen Rasmussen

17 *The Asthma and Allergy Association, Universitetsparken 4, 4000 Roskilde, Denmark*

18

19 Ole Hertel

20 *Department of Environmental Science – Atmospheric chemistry and physics, Aarhus University, Frederiksborgvej 399,*

21 *4000 Roskilde, Denmark*

22

23 **Abstract:** *Cladosporium* spp. are omnipresent moulds that grow on multiple substrates. Their spores possess a high
24 allergenic potential. Currently, little is known about the incidence and the sources of airborne *Cladosporium* spores in
25 Denmark. Air samples were collected between 31 May and 22 September 2015 in Viborg (Jutland, western Denmark).
26 Eighteen out of 21 days with daily average concentrations exceeding the health relevant threshold of 3,000 Spores m⁻³,
27 including the day with peak daily (13,553 Spores m⁻³) and 3-h concentrations (35,662 Spores m⁻³), occurred in August.
28 The air masses that approached Viborg during the longest episode of elevated spore concentrations originated from
29 northern Poland, the Baltics, passing over southern Sweden and the eastern Danish island of Zealand. The *Cladosporium*
30 spore concentrations from Viborg were compared with the *Cladosporium* spore concentrations from the operational
31 monitoring station in Copenhagen (Zealand, eastern Denmark). During the episode concentrations in Viborg were on
32 average 2,268 spores m⁻³ higher than in Copenhagen. On the peak day between 8:00-15:00 concentrations in Viborg
33 were 4-7 times higher than in Copenhagen, which we associated with grain crop harvesting in eastern Jutland. Elevated
34 day time concentrations in Viborg on the days with daily average concentrations exceeding the threshold also indicate the
35 local character of the sources.

36

37 **Keywords:** Aerobiology; *Cladosporium* spp.; Back Trajectories; HYSPLIT; Airmass Transport; Grain harvesting.

38

39 *Corresponding author e-mail: yo@ph.au.dk

40

41 **1 Introduction**

42 In many areas conidia of genus *Cladosporium* can account for more than 80% of all measured air spora. Spores
43 of *Cladosporium* spp. are well known aeroallergens, that affect the severity of asthma and allergic rhinitis in sensitized
44 persons (Denning et al. 2006). *Cladosporium* spp. grow on multiple substrates but mainly on decaying organic material,
45 which complicates assigning them to specific sources (Awad 2005). Additionally, due to their small size, these spores
46 can remain airborne for days while being transported from the source over great distances by wind in the absence of
47 precipitation. Identifying the geographical source area assists in determining the possible sources of *Cladosporium*
48 spores. The biological cycles and physical treatments, as e.g. agricultural operations, typical for the established sources,
49 can provide an important insight on the expected time window for intensive spore release, which can aid in managing
50 allergy outbreaks.

51 **2 Materials and methods**

52 Airborne spores of *Cladosporium* spp. were sampled by the Asthma and Allergy Association on the roof of
53 Regionshospitalet, 21 m above sea level in Viborg (56° 27'N 9° 24'E) during 115 days, 31 May–22 September 2015, on
54 the 48x14mm slides using a Hirst-type spore trap (Hirst 1952). The spores were counted at the genus level along 8
55 transversal transects at 6 mm distance between each under 630x magnification with further conversion to the spore air
56 concentrations with time resolution of 3h. Spore Integrals (SIns) were calculated by summing the average daily
57 concentrations over the specified periods (Galan et al. 2017).

58 The longest period of days (14–25 August) with daily average concentrations above the health relevant
59 threshold of 3,000 Spores m⁻³ (Gravesen 1979), containing the peak daily (Fig.1) and 3-h concentrations (Fig.2a),
60 together with the preceding day (13 August) and the following day (25 August) were selected for a detailed analysis of
61 air mass transport by back trajectories (Fig.2b). The back trajectories were computed by the HYSPLIT model (Draxler
62 and Hess 1998) with the Global Data Analysis System (GDAS) meteorological files in the form of a 1° latitude-longitude
63 grid with a receiving height of 500m and plotted 48h back in time with 3-h interval. The character of air mass transport
64 on these days presented an opportunity to compare 3-h *Cladosporium* spp. concentrations at the Viborg station and at the
65 Copenhagen station, the latter provided by the Asthma and Allergy Association. The description of the Copenhagen
66 station can be found elsewhere (Skjøth et al. 2012). The maps of grain agricultural fields (Fig.2c, Fig.2d) were created in
67 QGIS ver. 2.18 (<https://qgis.org/en/site/>).

68 The intra-diurnal *Cladosporium* spore cycles were plotted for every 3h of day as an average of corresponding
69 concentrations at the Viborg station.

70 Daily precipitation, daily averages of temperature (T_{mean}) and relative humidity (RH) were measured at the
71 meteorological station in Foulum (56° 29'N 9° 34'E).

72 **3 Results**

73 SIn over the study period totalled 218,151 Spore day m⁻³ with the average $T_{\text{mean}} = 14.3$ °C and RH = 79.5%. The
74 largest rainfall was observed in June (109.5mm), followed by July (72.5mm), September (64.4mm) and August (55 mm).
75 August was the warmest month ($T_{\text{mean}} = 16.9$ °C), whereas June was the coldest ($T_{\text{mean}} = 12$ °C). A positive statistically
76 significant relationship with daily spore concentrations was found only with T_{mean} ($r = 0.69$), whereas insignificant close to
77 zero spearman correlation coefficients were found for daily precipitation ($r = -0.03$) and RH ($r = -0.06$). The highest spore
78 concentrations were measured in August, which monthly SIn constituted 59.21% of the study SIn and equalled to
79 129,171 Spore day m⁻³, while July and September SIns were around 40,000 Spore day m⁻³ each.
80 Eighteen (out of 21) days with daily average concentration above 3,000 Spores m⁻³ occurred in August, contributing up

81 to 80% of August SIn. The peak daily average and 3-h concentrations were measured on 16 August (Fig.1; Fig.2A).

82 During the 11 days, 14–24 August, the air masses were arriving from the East and South East, i.e. originating in
83 the areas in Poland, the Baltic countries, and north-west Russia, passing over the Baltic sea and southern Sweden
84 (Fig.2b). Conversely, on 13 August and on 25 August the wind directions were distinctively different with the air masses
85 arriving from the North- and South-West (Fig. 2b). In the course of those 11 days the corresponding daily average
86 concentrations at the Copenhagen station were lower than at the Viborg station, with the exception of 15 August, when
87 the daily spore concentration in Viborg dropped down to 1,979 Spores m⁻³ while precipitation increased to 12.1 mm. On
88 16 August the daily average concentration at the Copenhagen station was less than half (5,685 Spores m⁻³) of the
89 concentration at the Viborg station (13,553 Spores m⁻³).

90 Maps of possible agricultural sources associated with the increase in *Cladosporium* spore air concentrations
91 during 14-24 August are shown in Fig.2c-d. In the area of eastern Jutland agricultural fields with winter grain seeds were
92 the main agricultural land cover (Fig.2c), with winter wheat as the dominating type of crop (Fig.2d).

93 The diurnal cycle of *Cladosporium* spores (Fig.3) on the days with daily average concentrations above 3,000
94 Spores m⁻³ had a maximum between 08:00 and 10:00. However, after excluding the high concentrations on 14 August
95 and 16 August, the diurnal distribution reflected elevated concentrations between 06:00 and 20:00 (Fig.3). No clear
96 diurnal pattern was found for the other 94 days with daily average concentrations below the threshold (Fig.3).

97 **4 Discussion**

98 *Cladosporium* conidia belong to the so-called dry spores, as higher temperatures and absence of precipitation
99 have been frequently observed to facilitate the increase in their air concentrations (Kasprzyk et al. 2016; Aira et al.
100 2012). The value of spearman correlation coefficient for the T_{mean} found in this study is comparable to the reported
101 previously, and despite the absence of correlation with precipitation, the highest concentrations of spores were observed
102 on the dry days.

103 The monthly SIns, the number of days with average concentration above 3,000 Spores m⁻³, and the value of the
104 peak concentrations in Viborg were lower compared with the values reported from Poland (Kasprzyk et al. 2016; Grinn-
105 Gofroń and Mika 2008), England (Sadyś et al. 2016), and France (Sindt et al. 2016) but higher than those found in
106 Morocco (Bardei et al. 2017), Croatia (Peternel et al. 2003) and northern Portugal (Oliveira et al. 2009). The climatic
107 conditions and the type of land cover determine both the availability of organic material in the form of vegetation, that is
108 prerequisite nutrient for fungi, and the growth of fungi mycelium leading to sporulation. Therefore, *Cladosporium* air
109 concentrations vary between different locations and between different years at the same location. We found August to be
110 the most important month in terms of airborne *Cladosporium* spore concentrations in 2015. This mono-modal monthly
111 distribution could reflect a typical pattern for *Cladosporium* in Viborg. However, in 2015 the T_{mean} in July was 2 degrees
112 lower than in August. This is unusual for Denmark, as July and August are normally the warmest months of the year with
113 ca. equal monthly T_{mean} values (www.dmi.dk). The daily pattern with higher concentrations between 08:00 and 17:00
114 based on the 19 days when *Cladosporium* concentrations were exceeding the health relevant threshold in Viborg (Fig.3)
115 indicates a local character of the sources rather than a distant.

116 On the peak day (16 August) the air masses originated in Latvia and western Belarus, passing over Lithuania,
117 north-western Poland, southern Sweden, and northern Zealand including Copenhagen (Fig.2b). Long distance transport
118 of air masses originating from the sources in Poland, Germany, and southern Sweden has been shown to affect air
119 concentrations of birch pollen and *Alternaria* spores in Copenhagen (Skjøth et al. 2012; Skjøth et al. 2007). Similarly,
120 the influence of the distant sources in southern Sweden, the Baltics, and Poland on *Cladosporium* air concentrations in

121 Viborg was visible in this study during the longest period of elevated concentrations (14–24 August) (Fig.2b). The
122 coastal position of Copenhagen is exceptional in the sense that its urbanised territory extends 20-30 km to the West and
123 North, whereas the East and South of the city are surrounded by the sea. Thus, in case of the wind coming from the East
124 and South, high concentrations of aerobiological particles in Copenhagen would be mainly influenced by the sources
125 located overseas. Viborg, unlike Copenhagen, is located in the centre of the Jutland peninsula and separated from the east
126 coast by ca. 100 km of mainly agricultural land. Therefore, the concentrations in Viborg which were 4-7 times higher
127 than in Copenhagen on 16 August (between 08:00-15:00) may have been supplemented by the yield of the sources
128 situated in closer proximity, i.e. in eastern Jutland. Additionally, during 14–24 August the daily *Cladosporium*
129 concentrations in Viborg were on average 2,268 Spores m⁻³ higher than in Copenhagen, which also indicates the
130 influence of local sources. According to the annual Danish report by the Agricultural Agency (Pedersen 2015), 14-24
131 August coincided with spring and winter grain crops harvesting activities.

132 This study provides the evidence that distant sources located in the Baltics, northern Poland, and southern
133 Sweden can affect air concentrations of *Cladosporium* in both eastern and western Denmark. However, harvesting of
134 grain crops, such as winter wheat, spring barley, winter rapeseed, barley, rye, winter and spring oats may have
135 contributed to the peak concentrations of *Cladosporium* in Viborg. A longer time-series in the eastern part of Denmark
136 will help to estimate the yield of local sources to peak *Cladosporium* concentrations in the area and to establish
137 *Cladosporium* monthly distribution and season duration. Additionally, the routine monitoring of spore concentrations in
138 Viborg can improve the accuracy of public information on *Cladosporium* concentrations, which is highly relevant for
139 sensitized individuals.

140

141 Acknowledgments

142 Data on agricultural grain fields land cover were provided by the Agricultural Agency under the Ministry of Environment
143 and Food of Denmark. Meteorological data were provided by the Danish Meteorological Institute.

144

145 References

- 146
147 Aira, M.-J., Rodríguez-Rajo, F.-J., Fernández-González, M., Seijo, C., Elvira-Rendueles, B., Gutiérrez-Bustillo, M., et al.
148 (2012). *Cladosporium* airborne spore incidence in the environmental quality of the Iberian Peninsula. *Grana*,
149 51(4), 293-304.
- 150 Awad, A. H. A. (2005). Vegetation: A source of air fungal bio-contaminant. *Aerobiologia*, 21(1), 53-61.
- 151 Bardei, F., Bouziane, H., Trigo, M. d. M., Ajouray, N., El Haskouri, F., & Kadiri, M. (2017). Atmospheric
152 concentrations and intradiurnal pattern of *Alternaria* and *Cladosporium* conidia in Tétouan (NW of Morocco).
153 *Aerobiologia*, 33(2), 221-228.
- 154 Denning, D. W., O'Driscoll, B. R., Hogaboam, C. M., Bowyer, P., & Niven, R. M. (2006). The link between fungi and
155 severe asthma: a summary of the evidence. *European Respiratory Journal*, 27(3), 615-626.
- 156 Draxler, R. R., & Hess, G. D. (1998). An overview of the HYSPLIT_4 modelling system for trajectories, dispersion and
157 deposition. *Australian meteorological magazine*, 47, 295-308.
- 158 Galan, C., Ariatti, A., Bonini, M., Clot, B., Crouzy, B., Dahl, A., et al. (2017). Recommended terminology for
159 aerobiological studies. *Aerbiologia*, 33, 293-295.
- 160 Gravesen, S. (1979). Fungi as a cause of allergic disease. *Allergy*, 34(3), 135-154.
- 161 Grinn-Gofroń, A., & Mika, A. (2008). Selected airborne allergenic fungal spores and meteorological factors in Szczecin,
162 Poland, 2004–2006. *Aerobiologia*, 24(2), 89.
- 163 Hirst, J. M. (1952). An automatic volumetric spore trap. *Annals of Applied Biology*, 39(2), 257-265.
- 164 Kasprzyk, I., Kaszewski, B. M., Weryszko-Chmielewska, E., Nowak, M., Sulborska, A., Kaczmarek, J., et al. (2016).
165 Warm and dry weather accelerates and elongates *Cladosporium* spore seasons in Poland. *Aerobiologia*, 32(1),
166 109-126.
- 167 Oliveira, M., Ribeiro, H., Delgado, J. L., & Abreu, I. (2009). Seasonal and intradiurnal variation of allergenic fungal
168 spores in urban and rural areas of the North of Portugal. *Aerobiologia*, 25(2), 85-98.

- 169 Pedersen, C. Å. (2015). Forsøgsarbejdet og vækstvilkår. In J. B. Pedersen (Ed.), *Oversigt over landsforsøgene 2015.*
170 *Forsøg og undersøgelser i Dansk Landbrugsrådgivning.* Aarhus, Denmark: SEGES P/S.
- 171 Peternel, R., Culig, J., & Hrga, I. (2003). Atmospheric concentrations of Cladosporium spp. and Alternaria spp. spores in
172 Zagreb (Croatia) and effects of some meteorological factors. *Annals of agricultural and environmental*
173 *medicine: AAEM, 11(2), 303-307.*
- 174 Sadyś, M., Adams-Groom, B., Herbert, R. J., & Kennedy, R. (2016). Comparisons of fungal spore distributions using air
175 sampling at Worcester, England (2006–2010). *Aerobiologia, 32(4), 619-634.*
- 176 Sindt, C., Besancenot, J.-P., & Thibaudon, M. (2016). Airborne Cladosporium fungal spores and climate change in
177 France. *Aerobiologia, 32(1), 53-68.*
- 178 Skjøth, C. A., Sommer, J., Frederiksen, L., & Gosewinkel Karlson, U. (2012). Crop harvest in Denmark and Central
179 Europe contributes to the local load of airborne Alternaria spore concentrations in Copenhagen. *Atmospheric*
180 *Chemistry and Physics, 12(22), 11107-11123.*
- 181 Skjøth, C. A., Sommer, J., Stach, A., Smith, M., & Brandt, J. (2007). The long-range transport of birch (Betula) pollen
182 from Poland and Germany causes significant pre-season concentrations in Denmark. *Clinical & Experimental*
183 *Allergy, 37(8), 1204-1212.*
- 184