

**Original citation:**

Alex J. Wadley, Gary Keane, Tom Cullen, Lynsey James, Jordan Vautrinot, Matthew Davies, Bethan Hussey, David J. Hunter, Sarabjit S. Mastana, Adrian Holliday, Steen Vang Petersen, Nicolette C. Bishop, Martin R. Lindley, and Steven J. Coles (2019) Characterisation of extracellular redox enzyme concentrations in response to exercise in humans, *Journal of Applied Physiology*, DOI: 10.1152/jappphysiol.00340.2019

**Permanent WRaP URL:**

<https://eprints.worc.ac.uk/8271/>

**Copyright and reuse:**

The Worcester Research and Publications (WRaP) makes this work available open access under the following conditions. Copyright © and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable the material made available in WRaP has been checked for eligibility before being made available.

Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

**Publisher's statement:**

© 2019 the American Physiological Society. This is an Accepted Manuscript of an article published online by the *Journal of Applied Physiology* on 27 June 2019, available online:

<https://doi.org/10.1152/jappphysiol.00340.2019>

**A note on versions:**

The version presented here may differ from the published version or, version of record, if you wish to cite this item you are advised to consult the publisher's version. Please see the 'permanent WRaP URL' above for details on accessing the published version and note that access may require a subscription.

**For more information, please contact [wrapteam@worc.ac.uk](mailto:wrapteam@worc.ac.uk)**

1 **Characterisation of extracellular redox enzyme concentrations in response**  
2 **to exercise in humans**

3 *Alex, J. Wadley<sup>a,b</sup> Gary Keane<sup>d</sup> Tom Cullen<sup>e</sup> Lynsey James<sup>a,c</sup> Jordan Vautrinot<sup>d</sup> Matthew*  
4 *Davies<sup>f</sup> Bethan Hussey<sup>a,c</sup> David, J. Hunter<sup>a,c</sup> Sarabjit Mastana<sup>a,c</sup> Adrian Holliday<sup>g</sup> Steen*  
5 *V. Petersen<sup>h</sup> Nicolette, C. Bishop<sup>a,b</sup> Martin, R. Lindley<sup>a,c\*</sup> & Steven, J. Coles<sup>d\*</sup>*

6  
7 <sup>a</sup> National Centre for Sport and Exercise Medicine, School of Sport, Exercise and Health  
8 Sciences, Loughborough University, Epinal Way, Loughborough LE11 3TU

9 <sup>b</sup> University Hospitals of Leicester NHS Trust, Infirmary Square, Leicester LE1 5WW

10 <sup>c</sup> Translational Chemical Biology Research Group, School of Sport, Exercise and Health  
11 Sciences, Loughborough University, LE11 3TU

12 <sup>d</sup> Institute of Science & the Environment, University of Worcester, Worcestershire, WR2 6AJ

13 <sup>e</sup> Centre for Sport, Exercise and Life Sciences, Coventry University, Coventry, CV15FB

14 <sup>f</sup> Institute of Sport & Exercise Sciences, University of Worcester, Worcestershire, WR2 6AJ

15 <sup>g</sup> Institute of Sport, Physical Activity and Leisure, Leeds Beckett University, LS1 3HE

16 <sup>h</sup> Department of Biomedicine, Aarhus University, DK-8000 Aarhus, Denmark

17 \* Both authors contributed equally

18  
19 **Keywords:** Antioxidant, oxidative stress, redoxkine, reactive oxygen species, peroxiredoxin

20  
21 **Running Title:** Extracellular redox enzymes & exercise

22  
23 **Word Count:** 4206

24  
25 **Corresponding author**

26 Dr Alex J Wadley

27 National Centre for Sport and Exercise Medicine

28 School of Sport, Exercise & Health Sciences

29 Loughborough University

30 Loughborough, UK, LE11 3TU

31 Email: a.j.wadley@lboro.ac.uk

32 Phone: 01509 223497

35 **Abstract**

36 Redox enzymes modulate intracellular redox balance and are secreted in response to  
37 cellular oxidative stress, potentially modulating systemic inflammation. Both aerobic and  
38 resistance exercise are known to cause acute systemic oxidative stress and inflammation;  
39 however, how redox enzyme concentrations alter in extracellular fluids following bouts of  
40 either type of exercise is unknown. Recreationally active males (n=26, mean  $\pm$  SD: age 28  $\pm$   
41 8 years) took part in either: 1) two separate energy-matched cycling bouts: one of moderate  
42 intensity (MOD) and a bout of high intensity interval exercise (HIIE) or 2) an eccentric-based  
43 resistance exercise protocol (RES). Alterations in plasma (study 1) and serum (study 2)  
44 peroxiredoxin (PRDX)-2, PRDX-4, superoxide dismutase-3 (SOD3), thioredoxin (TRX-1),  
45 TRX-reductase and interleukin (IL)-6 were assessed before and at various timepoints after  
46 exercise. There was a significant increase in SOD3 (+1.5 ng/mL) and PRDX-4 (+5.9 ng/mL)  
47 concentration following HIIE only, peaking at 30- and 60-min post-exercise respectively.  
48 TRX-R decreased immediately and 60-min following HIIE (-7.3 ng/mL) and MOD (-8.6  
49 ng/mL) respectively. In non-resistance trained males, no significant changes in redox enzyme  
50 concentrations were observed up to 48 hours following RES, despite significant muscle  
51 damage. IL-6 concentration increased in response to all trials, however there was no  
52 significant relationship between absolute or exercise-induced changes in redox enzyme  
53 concentrations. These results collectively suggest that HIIE, but not MOD or RES increase  
54 the extracellular concentration of PRDX-4 and SOD3. Exercise-induced changes in redox  
55 enzyme concentrations do not appear to directly relate to systemic changes in IL-6  
56 concentration.

57

58 **Abbreviations:** ANOVA: Analysis of Variance, BMI: Body Mass Index, CK: Creatine  
59 Kinase, ELISA: Enzyme Linked Immunosorbent Assay, EV: Extracellular Vesicle, H<sub>2</sub>O<sub>2</sub>:  
60 Hydrogen Peroxide, HIIE: High Intensity Interval Exercise, IL: Interleukin, IPAQ:  
61 International Physical Activity Questionnaire, LDH: Lactate Dehydrogenase, MOD:  
62 Moderate Intensity Exercise, NADH: reduced nicotinamide adenine dinucleotide, ONOO<sup>-</sup>:  
63 Peroxynitrite, PBS: Phosphate Buffered Saline, PBSwC: Phosphate Buffered Saline Wash  
64 Casein, PRDX: Peroxiredoxin, ROS: Reactive oxygen species, SD: Standard deviation, SOD:  
65 Superoxide Dismutase, TLR: Toll-like Receptor, TRX: Thioredoxin, TRX-R: Thioredoxin-  
66 Reductase, VO<sub>2MAX</sub>: Maximum oxygen consumption.

67 **New & Noteworthy**

68 We conducted two studies to characterise changes in redox enzyme concentrations after  
69 single bouts of exercise to investigate the emerging association between extracellular redox  
70 enzymes and inflammation. We provide evidence that SOD3 and PRDX-4 concentration  
71 increased following high intensity aerobic, but not eccentric-based resistance exercise.  
72 Changes were not associated with IL-6. The results provide a platform to investigate the  
73 utility of SOD3 and PRDX-4 as biomarkers of oxidative stress following exercise.

74

75

76

77

78

79

80

81

82

83

84

85

86

87

88

89

90

91

92 **Introduction**

93           It is well documented that acute exercise perturbs cellular reduction-oxidation (redox)  
94 balance through the increased production of reactive oxygen species (ROS) within actively  
95 contracting skeletal muscle (34), as well as other infiltrating cell types (35). Evidence  
96 suggests that ROS such as hydrogen peroxide ( $H_2O_2$ ) and peroxynitrite ( $ONOO^-$ ) have  
97 important roles in facilitating muscle contractile activity (25) and regulating the expression of  
98 genes involved with metabolism and endogenous antioxidant protection (14, 39). Conversely,  
99 heightened levels of exercise-induced  $H_2O_2$  at the expense of antioxidant defense systems can  
100 elicit oxidative stress, which may limit contractile function and promote fatigue (33). Given  
101 this biphasic relationship, studies have previously evaluated alterations in redox balance in  
102 response to both aerobic and resistance type exercise. These studies have primarily focused  
103 on the quantification of distal markers in extracellular fluids, such as the oxidation  
104 biomolecules and/or activity of antioxidant enzymes in plasma (48), serum (31), saliva (11)  
105 and urine (41); highlighting exercise duration (3), intensity (17) and muscle-damage (4) as  
106 factors governing greater increases. However, criticisms are commonly made with regards to  
107 the direct relationship of these markers with the redox state of active tissues during exercise  
108 (9). Recent evidence has highlighted that intracellular redox enzymes, such as peroxiredoxin  
109 (PRDX) can be secreted from skeletal muscle myocytes (28) and immune cells (40) in  
110 response to increasing concentrations of  $H_2O_2$  *in vitro*. Human studies are also beginning to  
111 provide evidence that plasma/ serum PRDX-2 and PRDX-4 concentrations **could** serve as  
112 important biomarkers of intracellular redox state in the context of acute and chronic  
113 inflammatory conditions (27, 40).

114           PRDXs are a major family of ubiquitous redox proteins, which modulate intracellular  
115 redox balance through a highly reactive cysteine thiolate group. The reaction rate of this  
116 cysteine is markedly greater than any other thiol-containing protein (50), allowing rapid  
117 regulation of cellular  $H_2O_2$ , with some evidence to suggest that this may facilitate muscle  
118 contraction (26). PRDXs are therefore reliable footprints of intracellular redox state, with  
119 heightened oxidation of the PRDX cysteine indicative of oxidative stress (37). In addition,  
120 upon secretion from immune cells, PRDX can directly bind to toll-like receptor (TLR)-4 to  
121 initiate inflammatory cytokine production (e.g. interleukin (IL)-6) (38), providing some  
122 support for the association between PRDX and inflammation (27, 40). Recent work has  
123 begun to explore changes in the PRDX catalytic cycle in blood cells isolated from humans  
124 before and after acute exercise (6, 46, 47). In parallel with increases in soluble markers of  
125 inflammation (e.g. IL-6 and C-reactive protein), an increase in the oxidation of PRDX (i.e.

126 dimer and over-oxidised states) has been reported following intensive cycling and running  
127 exercise (46, 47). To our knowledge, changes in PRDX have yet to be assessed in the context  
128 of exercise in humans and represents a potentially unexplored area of exercise and redox  
129 biology. Interestingly, PRDX-2 can be secreted in tandem with its enzymatic reducing  
130 partners, thioredoxin (TRX-1) and thioredoxin reductase (TRX-R) (20, 40). TRX-1 and TRX-  
131 R are cysteine and selenium based-antioxidant enzymes respectively, with higher reduction  
132 potentials than PRDX, thus contributing towards maintaining the antioxidant function of  
133 PRDX. In addition, the enzyme superoxide dismutase 3 (SOD3) is an extracellular  
134 antioxidant released upon cellular stimulation, providing an immediate change in  
135 extracellular antioxidant capacity (15, 20). Given the emerging body of literature supporting a  
136 relationship between intracellular oxidative stress, redox enzyme secretion and soluble  
137 inflammatory markers, the quantification of PRDX-2, PRDX-4, TRX-1, TRX-R and SOD3 in  
138 extracellular fluids offers the potential for accurate assessment of changes in oxidative stress  
139 and inflammation after different types of exercise.

140         Based upon existing knowledge of the factors that can impact acute changes in  
141 exercise-induced oxidative stress, we sought to perform two experiments to understand how  
142 novel markers, such as PRDX-2, PRDX-4, TRX-1, TRX-R and SOD3 respond to acute  
143 exercise, and whether relationships exist between changes in inflammation. Specifically, we  
144 aimed to characterise how these markers would be impacted by aerobic exercise intensity and  
145 eccentric-based resistance exercise. We tested the hypothesis that both protocols would elicit  
146 an increase in the concentrations of redox enzymes within plasma/ serum after exercise; with  
147 higher exercise intensity causing a larger increase following aerobic exercise.

148

## 149 **Methods**

### 150 Participants

151         Healthy, untrained participants were recruited for two independent studies (Table 1)  
152 Participants in both studies completed the International Physical Activity Questionnaire  
153 (IPAQ), which addresses habitual levels of weekly physical activity. Participants gave their  
154 informed written consent and all studies were approved by the local Ethical Review  
155 Committee, in accordance with the Declaration of Helsinki, 2008. Participants were all non-  
156 smokers and had not taken any antioxidant vitamin supplements or anti-inflammatory drugs  
157 for 8 weeks prior to the laboratory visits. All participants were required to refrain from any

158 strenuous physical activity, consumption of alcoholic beverages or caffeine for at least two  
159 days prior to the experimental sessions.

160

### 161 Experimental Sessions

162 The full workflow for this project is detailed in Figure 1. Experimental sessions took  
163 place in the morning (7.00 - 8.00 am start time) under stable climatic conditions (18 - 20°C  
164 and humidity between 45 – 55%) and following at least a 10-hour fast. After a period of rest,  
165 height (*Seca Alpha, Hamburg, Germany*) and mass (*Tanita, Tokyo, Japan*) were determined.

166 In study 1, participants first visited the laboratory for an assessment of  
167 cardiorespiratory fitness ( $\dot{V}O_{2\text{MAX}}$ ) using a ramp test to exhaustion on an electromagnetically  
168 braked cycle ergometer (*Lode Excalibur Sport, Groningen, Netherlands*). The protocol  
169 involved commencing pedalling at 100 Watts, followed by fixed 30-Watt increments every 4  
170 minutes. Oxygen uptake was assessed continuously using a breath-by-breath system (*Oxycon*  
171 *Pro, Jaeger, Wuertzberg, Germany*) and heart rate monitored using a Polar Vantage heart rate  
172 monitor (*Polar, Kempele, Finland*). The test ended when the participant reached volitional  
173 exhaustion or when a plateau in oxygen consumption was observed with an increase in  
174 workload (49). A final obtained value of rate of oxygen consumption was accepted as  $\dot{V}O_{2\text{MAX}}$   
175 and expressed relative to body weight ( $\text{mL}\cdot\text{kg}^{-1}\cdot\text{min}^{-1}$ ). At least one week later,  
176 participants then undertook the first of two energy and time-matched cycling trials in a  
177 randomised order, at least one week apart: a continuous bout of moderate intensity cycling at  
178 approximately 60%  $\dot{V}O_{2\text{MAX}}$  for 58 minutes (MOD) and a bout of high intensity interval  
179 exercise (HIIE), consisting of 10 x 4-minute intervals at 85%  $\dot{V}O_{2\text{MAX}}$ , with 2-minute rest  
180 intervals. In both trials, oxygen uptake was assessed continuously and power output was  
181 adjusted where necessary in order to maintain target  $\dot{V}O_{2\text{MAX}}$  and equal energy expenditure  
182 between MOD and HIIE (study 1). Rating of Perceived Exertion (RPE) was monitored every  
183 5 minutes throughout the trials (5).

184 In study 2 (n = 16), non-resistance trained males undertook an eccentric-based  
185 resistance exercise protocol adapted from a previous study by Alemany et al (1). This muscle  
186 damaging protocol was performed on a Humac Norm dynamometer (CSMI, Massachusetts,  
187 USA). The dynamometer lever arm was programmed to flex the participant's knee from a  
188 start position of 10° of flexion to 90° of flexion, thus allowing a range of motion of 80°. The  
189 participants began with their leg at the start position and were asked to maximally contract  
190 their quadriceps against a resistance while the lever arm moved to the finish position (90°

191 knee flexion). Once at the finish position they were advised to relax their leg and the  
192 dynamometer moved them back to the start position to avoid a concentric contraction being  
193 performed. The lever arm moved at a set speed of  $60^{\circ}\cdot\text{s}^{-1}$ . The bout consisted of 20 sets of 10  
194 repetitions with each set being separated by 1 minute **of** rest. Visual feedback and verbal  
195 encouragement **were** provided to all participants to maximise torque output for each  
196 contraction.

197

### 198 Blood sampling and Plasma Isolation

199 For both studies, a catheter (*Appleton Woods, Birmingham, UK*) was inserted into the  
200 antecubital vein of the arm prior to exercise to obtain a baseline blood sample after thirty  
201 minutes of rest (Pre). The catheter was continually kept clear with isotonic saline solution  
202 (0.9% sodium chloride). As indicated in Figure 1, blood samples were then taken  
203 immediately, 30 minutes and 60 minutes after both HIIE and MOD (Study 1 – Pre, Post+0,  
204 Post+30 and Post+60) and immediately, 30 minutes, 3 hours and 48 hours following the  
205 muscle damage protocol (Study 2 – Pre, Post-0, Post-30min, Post+3hr and Post+48hr). The  
206 post+48 hr (Study 2) blood sample was taken via venepuncture. At each time point, 12 mL of  
207 blood was drawn into vacutainer tubes containing either potassium ethylene  
208 diaminetetraacetic acid in study 1 (*Becton, Dickson & Company, Oxford, UK*) or no  
209 anticoagulant in study 2. In study 1, whole blood was centrifuged at 1525g for 15 minutes, at  
210 room temperature. In study 2, whole blood was allowed to clot at room temperature for 20  
211 mins and then centrifuged at 1500g for 15 minutes. The resulting plasma (study 1) and serum  
212 (study 2) were aliquoted and frozen at  $-80^{\circ}\text{C}$  for future analysis of redox enzymes, IL-6,  
213 creatine kinase (CK) and lactate dehydrogenase (LDH). Capillary blood samples were  
214 obtained from the earlobe after 4 min of exercise and then every 6 min thereafter (i.e. end of  
215 each HIIE interval) in study 1. These samples were used for analysis of blood glucose and  
216 lactate concentrations to verify intensity-dependent differences between each protocol.

217

### 218 Analytical Procedures

#### 219 *PRDX-2, PRDX-4, TRX-1, TRX-R and SOD3 ELISAs*

220 ELISAs for the detection of PRDX-2, PRDX-4, TRX, TRX-R and SOD3 were  
221 developed in-house. Commercially available antigens and antibodies (i.e. PRDX-2, PRDX-4,  
222 TRX and TRX-R) were purchased from either *Abcam*, Cambridge, UK (ab) or *Sigma Aldrich*,  
223 Dorset, UK (SRP). The human SOD3 antigen and rabbit antiserum directed against human  
224 SOD3 were developed as previously described (16, 20). Plasma or serum and standards (100

225  $\mu\text{L}$ ) were loaded onto individual wells of an ELISA plate (Thermo Scientific F8 polysorp  
226 immune wells) and protein left to bind overnight at 4 °C. Wells were then pre-washed with  
227 PBS wash buffer, supplemented with 0.1% casein (PBSwC, 200  $\mu\text{L}$ ) and then blocked with  
228 1% casein in PBS (200  $\mu\text{L}$ ) for 30 minutes at room temperature, with gentle agitation. Anti-  
229 human rabbit antibodies for PRDX-2 (ab133481, 1:2000), PRDX-4 (ab59542, 1:2000) and  
230 SOD3 (in-house, 1:2000), and anti-human mouse antibodies for TRX-1 (ab16965, 1:8000)  
231 and TRX-R (ab16847, 1:1000) were then added to each well, diluted in PBSwC for 45  
232 minutes at room temperature. Following this, 100  $\mu\text{L}$  of anti-rabbit (1:5000) or anti-mouse  
233 (1:500) IgG Biotin antibodies in PBSwC, and streptavidin-horseradish peroxidase (1:2000 in  
234 PBSwC) were added separately to each well, both for 45 minutes, with gentle agitation.  
235 Between all stages, all wells were washed three times with PBSwC. Finally, 100  $\mu\text{L}$  of  
236 3,3',5,5'-tetramethylbenzidine (10ug) was added per well, and the plate left to develop in the  
237 dark for 15-25 minutes. Stop solution (1.5mM  $\text{H}_2\text{SO}_4$ , 50  $\mu\text{L}$ ) was then added to each well  
238 and absorption at 450nm subsequently evaluated by using a plate reader (Multiskan Ascent,  
239 Thermo Labsystems). Concentration of each antigen was then determined by comparing  
240 absorbance values of recombinant PRDX-2 (ab167977, *Abcam*), PRDX-4 (ab93947, *Abcam*),  
241 TRX-1 (ab51064, *Abcam*), TRX-R (SRP6081, *Sigma Aldrich*) and SOD3 (in-house) proteins  
242 (0-50 ng/mL). ELISA validation experiments showed no cross-reactivity of the PRDX-2,  
243 PRDX-4, TRX-1, TRX-R and SOD3 antibodies with the respective antigens, nor with serum  
244 albumin. All values were adjusted for plasma volume, according to previous methods (12).

245

#### 246 *Other Analyses*

247 In both studies, a cytometric bead array was used to quantify plasma (study 1) and  
248 serum (study 2) IL-6 concentrations on a BD C6 Accuri Flow Cytometer (*BD Biosciences*,  
249 *Berkshire*). In study 1, blood lactate and glucose concentrations were determined  
250 immediately following collection using an automated lactate and glucose analyser (Biosen C-  
251 Line Clinic, EKF-diagnostic GmbH, *Barleben, Germany*). In study 2, serum CK and LDH  
252 concentrations were determined to monitor muscle damage using an automated ABX Pentra  
253 400 system (*Horiba UK Ltd, UK*). Haematocrit and haemoglobin concentrations were used to  
254 ascertain plasma volume changes and make appropriate adjustments in plasma redox enzyme  
255 and IL-6 concentrations (Beckman Coulter, *London, UK*).

256

#### 257 **Statistical Analysis**

258 The Shapiro Wilk test was used to test for normality in scale data at all time points.  
259 Differences between participant characteristics and the physiological responses to exercise in  
260 both studies were assessed using unpaired samples T-tests or non-parametric Mann-Whitney  
261 U Tests. The influence of exercise on plasma/ serum PRDX-2, PRDX-4, SOD3, TRX-1,  
262 TRX-R and IL-6 concentration was assessed over time by repeated-measures analysis of  
263 variance (ANOVA) or non-parametric Wilcoxon signed rank tests, depending variable  
264 normality. Post hoc analysis of any significant effect of time or interaction effect (study 1;  
265 Group\*Time) was performed by a test of simple effects by pairwise comparisons, with  
266 Bonferroni correction. Effect sizes for main effects and interaction effects of ANOVA are  
267 presented as partial eta<sup>2</sup> ( $\eta^2_p$ ), using Cohen's definition of  $\eta^2_p$  of 0.01, 0.06 and 0.14 for  
268 'small', 'medium' and 'large' effects respectively (10). Pearson correlation and Spearman  
269 rank were used to assess the relationship between parametric and non-parametric data  
270 respectively. All values are presented as means  $\pm$  standard deviation or error (indicated  
271 throughout manuscript). Statistical significance was accepted at the  $p < .05$  level. Statistical  
272 analyses were performed using SPSS (*PASW Statistics, release 23.0, SPSS Inc., Chicago, IL,*  
273 *USA*).

274

## 275 **Results**

276 There was no significant difference in age or BMI between the participants taking part  
277 in the two studies, Participants in study 1 ( $p = 0.004$ ) had significantly higher self-reported  
278 physical activity than in study 2.

279

### 280 *Acute physiological responses to HIIE and MOD*

281 For study 1, the physiological responses during each exercise bout are reported in  
282 Table 2. Peak  $\dot{V}O_2$  and RPE were significantly greater in HIIE compared to MOD ( $p <$   
283  $0.00001$ ), but there were no statistically significant differences in mean  $\dot{V}O_2$  and energy  
284 expenditure. Whole blood lactate and glucose data are reported in Table 2. Mean lactate  
285 concentration was significantly higher during HIIE than MOD ( $p < 0.0001$ ), but there was no  
286 significant difference in average glucose concentration between trials.

287

### 288 *Effects of eccentric-based resistance exercise on muscle damage markers*

289 Changes in markers of muscle damage are reported in Table 3. A stepwise increase  
290 (Post+48hr > Post+3hr > Post+30min > Post+0 > Pre) in serum CK concentration was

291 observed over time, peaking above Pre at Post+48hr ( $p > 0.001$ ). Serum LDH concentration  
292 was elevated above Pre at all post-exercise timepoints ( $p < 0.05$ ), also increasing Post+3hr  
293 and Post+48hr, relative to Post+30min ( $p < 0.05$ ).

294

#### 295 *Effects of aerobic and eccentric-based resistance exercise on IL-6 concentration*

296 IL-6 data is presented in Figure 3. In study 1, plasma IL-6 increased in both trials  
297 (Time effect:  $F(3) = 15.5$ ,  $p < 0.0001$ ,  $\eta^2 = 0.66$ ), being elevated above resting values, both  
298 immediately ( $p = 0.004$ ) and Post+30 ( $p = 0.002$ ), but not Post+60 (Figure 3A). The  
299 magnitude of this increase was significantly greater Post-Ex in HIIE ( $p = 0.031$ ), than MOD  
300 (Time x Condition effect:  $F(3) = 7.0$ ,  $p < 0.001$ ,  $\eta^2 = 0.47$ ). IL-6 concentration decreased  
301 Post+30 ( $p = 0.004$ ) and Post+60 ( $p = 0.007$ ), relative to Post+0, and Post+60, relative to  
302 Post+30 ( $p = 0.026$ ) in HIIE only. In study 2 (Figure 3B), IL-6 concentration was  
303 significantly higher at all timepoints up to three hours, but not 48 hours after exercise,  
304 relative to Pre (Time effect:  $F(4) = 14.3$ ,  $p < 0.0001$ ,  $\eta^2 = 0.30$ ).

305

#### 306 *Effects of aerobic exercise on PRDX-2, PRDX-4, TRX-1, TRX-R and SOD3 concentration*

307 No differences were observed in resting concentrations of PRDX-2, PRDX-4, TRX-1,  
308 TRX-R or SOD3 when quantified in plasma and serum across all trials. Changes in plasma  
309 PRDX-2, PRDX-4, TRX-1, TRX-R and SOD3 in response to MOD and HIIE are reported in  
310 Figure 2A. There was a significant increase in plasma SOD3 (Trial x Time Effect:  $F(3,1) =$   
311  $5.3$ ,  $p = 0.028$ ,  $\eta^2 = 0.31$ ) and PRDX-4 following HIIE only (non-parametric tests: all  $p <$   
312  $0.05$ ). SOD3 concentration was elevated above pre-exercise values at all post-HIIE  
313 timepoints, peaking at Post+0 ( $p = 0.015$ ) and Post+30 ( $p = 0.013$ ), but only significantly  
314 higher than MOD at Post+30 ( $p = 0.05$ ). Plasma SOD3 concentration decreased relative to  
315 Post+30 at Post+60 ( $p = 0.013$ ). Relative to Pre, PRDX-4 concentration increased at Post+30  
316 ( $p = 0.015$ ) and Post+60 ( $p = 0.008$ ) following HIIE, with PRDX-4 concentration higher at all  
317 post-exercise timepoints compared with MOD ( $p < 0.038$ ). There was a significant decrease  
318 in plasma TRX-R concentration in both MOD and HIIE. Relative to Pre, TRX-R significantly  
319 decreased at Post+0 in HIIE only ( $p = 0.021$ ), with values significantly less than MOD ( $p =$   
320  $0.011$ ). Following MOD, TRX-R was significantly lower at Post+60, relative to all  
321 timepoints (all  $p < 0.038$ ). There were no statistically significant changes in PRDX-2 and  
322 TRX-1 concentration over time in either trial; however, TRX-1 concentration was  
323 significantly higher in HIIE than MOD Post+60 only ( $p = 0.021$ ).

324

325 *Effects of eccentric-based resistance exercise on PRDX-2, PRDX-4, TRX-1, TRX-R and*  
326 *SOD3 concentration*

327 Serum redox enzyme concentration changes in response to an eccentric-based  
328 resistance exercise protocol are presented in Figure 2B. A trend was observed for a decrease  
329 in PRDX-2 concentration Post+30min (-1.12 ng/mL), however this did not reach statistical  
330 significance (Time effect:  $F(4) = 2.3$ ,  $p = 0.065$ ,  $\eta^2 = 0.13$ ). Similarly, no significant changes  
331 were noted in PRDX-4, TRX-R or SOD3 up to 48 hours following eccentric-based resistance  
332 exercise. A significant increase in TRX-1 was shown Post+48hr, relative to Post+30min ( $p =$   
333  $0.039$ ), but not Pre ( $p = 0.309$ ).

334

335

### 336 **Discussion**

337 The current results have characterised the kinetic responses of endogenous redox  
338 enzymes within the extracellular environment after exercise for the first time. We highlight  
339 novel findings that high intensity aerobic cycling induces a significant increase in SOD3 and  
340 PRDX-4 in healthy, untrained males. Similar responses were not observed following  
341 moderate intensity cycling or muscle damaging resistance exercise. In contrast, plasma TRX-  
342 R concentration decreased within one hour following moderate and high-intensity cycling  
343 exercise, but not resistance exercise. Taken together these findings provide novel insights into  
344 the regulation of extracellular redox enzymes in response to exercise.

345 The current data highlights modality and exercise-intensity specific increases in two  
346 abundant redox enzymes. In response to aerobic exercise, PRDX-4, but not PRDX-2  
347 concentration increased thirty minutes following HIIE and remained elevated until Post+60.  
348 The secretory pathways of PRDXs are isoform specific, with endoplasmic reticulum (ER, i.e.  
349 PRDX-4) and cytosolic (i.e. PRDX-2) resident isoforms released via classical and non-  
350 classical secretory pathways respectively (8). The current data therefore suggests that  
351 exercise may activate the ER-golgi pathway to secrete PRDX-4 in an intensity-dependent  
352 manner. SOD3, which is also released via this pathway, increased more rapidly than PRDX-4  
353 following HIIE (Post+0), with levels tailing off Post+60, relative to Post+30. SOD3 is an  
354 antioxidant enzyme released directly from the cell membrane (15, 20), specifically secreted  
355 during exercise to metabolise superoxide anions produced in the extracellular environment to  
356  $H_2O_2$  (30). The different peak concentrations of SOD3 (i.e. Post+0) and PRDX-4 (i.e.

357 Post+30) following HIIE may be explained, in part, by a) the membrane proximity of SOD3  
358 compared to the ER location of PRDX-4 and b) the appearance of superoxide anions first in  
359 the extracellular space following exercise, before their metabolism to H<sub>2</sub>O<sub>2</sub>, which then  
360 induced PRDX-4 secretion. This may also be reflective of differential secretion rates of  
361 SOD3 and PRDX-4 from various tissues during and following exercise. Both proteins are  
362 expressed in skeletal muscle (19), a highly redox active tissue (36); however, PRDX-4 is  
363 primarily located in pancreas, liver and heart (21), whereas SOD3 is expressed in the heart  
364 and vasculature tissue (42). The association with the vasculature may explain the more rapid  
365 increase in plasma SOD3 concentration following HIIE. Aside from these increases, a modest  
366 decrease was observed in plasma TRX-R after both MOD and HIIE (study 1), with this  
367 change being much more rapid in HIIE (Post+0), compared to MOD (Post+60). The  
368 mechanisms driving a decrease in TRX-R after exercise are unclear at present. The decrease  
369 may represent transient homeostatic fluctuations involving uptake of redox enzymes by  
370 neighboring cells and tissues, perhaps to regulate intracellular redox balance (23).

371 A finding that was in contrast to our hypothesis was that eccentric-based resistance  
372 exercise did not induce an increase in the extracellular concentrations of redox enzymes. The  
373 measurement of redox enzymes in plasma and serum is an emerging area of biomedical  
374 research, particularly in the context of acute (24) and chronic (13, 43) inflammatory  
375 conditions, where PRDXs and TRX-1 have been associated with enhanced cytokine and  
376 chemokine production (22, 38). The participants in both studies were relatively inactive, with  
377 participants in study 2 in particular, reporting significantly lower levels of habitual physical  
378 activity (Table 2) and being unaccustomed to eccentric-based resistance exercise.  
379 Unaccustomed eccentric exercise induces significant amounts of acute muscle damage and  
380 inflammation (7), as demonstrated by the stepwise increases in CK and LDH concentrations  
381 up to 48 hours following our protocol, and IL-6 up to 3 hours post-exercise (Figure 3B).  
382 These data suggest that the increase in SOD3 and PRDX-4 observed in study one is unlikely  
383 due to just a disruption to the plasma membrane, given that no changes were observed  
384 following a muscle-damaging bout of resistance exercise. It must be acknowledged that only  
385 selective timepoints were measured following the protocol, and perhaps the secretion of  
386 redox enzymes occurs between 3- and 48-hours post-exercise. Nevertheless, this study has  
387 highlighted for the first time that redox enzyme concentrations do not match that of  
388 established markers of muscle damage and inflammation when measured in serum samples  
389 following an eccentric-based resistance exercise bout. In response to aerobic-based exercise,

390 we have recently demonstrated a positive association between intracellular peroxiredoxin (I-  
391 IV) over-oxidation in immune cells and plasma IL-6 concentration (47). In the current study,  
392 IL-6 concentration increased in an intensity-dependent manner (HIIE > MOD) following  
393 aerobic exercise (Figure 3A); however, there were no statistically significant relationships  
394 between absolute or exercise-induced changes in PRDX-4 and SOD3 with IL-6. The  
395 observations across both studies therefore suggest no relationship between that IL-6 and  
396 redox enzymes after exercise. A larger sample size may be needed to adequately address  
397 these associations and support the previously documented relationship between plasma/  
398 serum redox enzymes and soluble inflammatory markers (27, 40).

399 The results of the current investigation demonstrate clear differences in the changes in  
400 SOD3, TRX-R and PRDX-4 following aerobic vs. eccentric-based resistance exercise. With  
401 regards to PRDX-2 and TRX-1, no changes were observed following aerobic or eccentric-  
402 based resistance exercise. Both PRDX-2 and TRX-1 are cytosolic redox enzymes that contain  
403 no N-terminal signal peptide for secretion and thus are released via non-classical pathways,  
404 associated with extracellular vesicles (EVs), such as exosomes and nanoparticles (45).  
405 PRDX-2 and TRX-1 are detectable in plasma/ serum samples through their association with  
406 the exofacial surface of the EV membrane (18, 44); however, their protein levels may be  
407 higher due to protein contained within the EVs. This protein would not be detectable by  
408 antibodies when enclosed within the lipid membrane during ELISA quantification, as  
409 previously shown (32). Indeed, recent evidence has highlighted that a series of leaderless  
410 redox enzymes (i.e, PRDX-1, PRDX-2, PRDX-5, PRDX-6, TRX-1, SOD1 and SOD2) are  
411 secreted in EVs via a non-classical route following exposure to stress, with classically  
412 secreted SOD3, TRX-R and PRDX-4 not detectable within EVs (2). This may explain why  
413 plasma/ serum PRDX-2 and TRX-1 concentration did not significantly change following  
414 muscle-damaging or aerobic exercise. It must be noted that TRX-1 concentration was  
415 significantly higher 48 hours after the eccentric-based resistance exercise protocol, relative to  
416 **Post+0** (study 2) and also significantly higher at Post+60 in HIIE, compared to MOD (study  
417 1). These findings again underpin intensity-dependent differences, despite in both cases,  
418 concentrations not being higher than pre-exercise values. In response to a far more extreme  
419 bout of exercise, Marumoto *et al*, (2010) reported a marked increase in TRX-1 levels  
420 ( $17.9 \pm 1.2$  ng/mL at baseline to  $70.1 \pm 6.9$  ng/mL) after a 2-day 130km ultra-endurance  
421 marathon (29); however, these exercise bouts were substantially different in nature and thus  
422 hard to directly compare. Even though an Ultramarathon is accompanied by significant

423 amounts of muscle damage, given the findings of study 2, it is unlikely that muscle damage is  
424 the primary cause of TRX-1 secretion in this context. Further work is needed to clarify  
425 whether TRX-1 and PRDX-2 protein levels alter within EVs after conventional bouts (i.e. not  
426 ultra-endurance) of muscle-damaging and aerobic-based exercise.

427 This study has quantified the responses of antioxidant enzymes in the extracellular  
428 environment following acute exercise in age and BMI matched individuals from two  
429 independent exercise studies (Table 1). We must acknowledge that the studies would have  
430 benefited from a direct comparison between redox enzyme concentrations and other  
431 established biomarkers of oxidative stress (e.g. protein carbonyls and F2-isoprostanes).  
432 However, due to limited sample volume this analysis was not feasible and should therefore be  
433 prioritised as an area of future research. A second limitation is that the quantification of redox  
434 enzymes and IL-6 were undertaken in both plasma (study 1) and serum (study 2); however,  
435 there were no differences in any of these proteins when quantified in pre-exercise samples.

436

## 437 **Conclusion**

438 The results of the present study have highlighted that plasma SOD3 and PRDX-4  
439 concentration increased in response to acute exercise. Importantly, the secretion of these  
440 proteins appears to be intensity and modality dependent, with increases only observed in  
441 response to high intensity aerobic cycling in untrained individuals. A decrease in TRX-R was  
442 also noted following different aerobic exercise bouts, with exercise intensity driving a more  
443 rapid decrease in TRX-R. Future research is required to pinpoint the precise mechanisms  
444 governing the secretion and uptake of redox enzymes, and their role in regulating redox  
445 balance between tissues after exercise.

446

447

448

449

## 450 **Acknowledgements**

451 This project involved analysis of blood samples from 2 independent exercise studies run by  
452 AW, TC, JV, GK, MD & SC (Study 1) and DH, BH, LJ, SM & ML (Study 2). AW and GK

453 carried out ELISA optimisation and subsequent analysis. AW wrote the manuscript and all  
454 authors commented to the final draft.

455

456 **Conflict of Interest**

457 None of the authors declare a conflict of interest.

458

459 **Funding**

460 This research was supported by the University of Worcester and the National Institute for  
461 Health Research (NIHR) Leicester Biomedical Research Centre. The views expressed are  
462 those of the authors and not necessarily those of the NHS, the NIHR or the Department of  
463 Health

## References

1. **Aleman JA, Mathews H, Davis JM, Kostek MC, States U, States U.** Comparison of Acute Responses to Isotonic or Isokinetic Eccentric Muscle Action : Differential Outcomes in Skeletal Muscle Damage and Implications for Rehabilitation. *Int J Sport Med* 35: 1–7, 2014.
2. **Bekeschus S, Moritz J, Schmidt A, Wende K.** Redox regulation of leukocyte-derived microparticle release and protein content in response to cold physical plasma-derived oxidants. *Clin Plasma Med* 7–8: 24–35, 2017.
3. **Bloomer RJ, Davis PG, Consitt LA, Wideman L.** Plasma protein carbonyl response to increasing exercise duration in aerobically trained men and women [Online]. *Int J Sports Med* 28: 21–25, 2007. <https://www.thieme-connect.de/ejournals/pdf/sportsmed/doi/10.1055/s-2006-924140.pdf>.
4. **Bloomer RJ, Goldfarb AH, Wideman L, McKenzie MJ, Consitt LA.** Effects of acute aerobic and anaerobic exercise on blood markers of oxidative stress. [Online]. *J Strength Cond Res* 19: 276–85, 2005. <http://www.ncbi.nlm.nih.gov/pubmed/15903362>.
5. **Borg G.** The psychophysical bases of perceived exertion. *Med Sci Sport Exerc* 14: 377–381, 1982.
6. **Brinkmann C, Blossfeld J, Pesch M, Krone B, Wiesiollek K, Capin D, Montiel G, Hellmich M, Bloch W, Brixius K.** Lipid-peroxidation and peroxiredoxin-overoxidation in the erythrocytes of non-insulin-dependent type 2 diabetic men during acute exercise. *Eur J Appl Physiol* 112: 2277–87, 2012.
7. **Bruunsgaard H, Galbo H, Halkjaer-Kristensen J, Johansen TL, MacLean DA, Pedersen BK.** Exercise-induced increase in serum interleukin-6 in humans is related to muscle damage. *J Physiol* 499: 833–841, 1997.
8. **Chang, J.W., Lee, S.H., Lu, Y., Yoo YJ.** Transforming growth factor- $\beta$ 1 induces the non-classical secretion of peroxiredoxin-I in A549 cells. *Biochem Biophys Res Commun* 345: 118–123, 2006.
9. **Cobley JN, Close GL, Bailey DM, Davison GW.** Exercise redox biochemistry: Conceptual, methodological and technical recommendations. *Redox Biol* 12: 540–548, 2017.

10. **Cohen J.** *Statistical Power Analysis for the Behavioural Sciences, 2nd Edition.* 1988.
11. **Deminice R, Sicchiere T, Payao PO, Jordao A.** Blood and salivary oxidative stress biomarker following an acute session of resistance exercise in humans. *Int J Sport Med* 31: 599–603, 2010.
12. **Dill DB, Costill DL.** Calculation of percentage changes in volumes of blood, plasma, and red cells in dehydration. *J Appl Physiol* 37: 247–248, 1974.
13. **Gerrits EG, Alkhalaf A, Landman GWD, van Hateren KJJ, Groenier KH, Struck J, Schulte J, Gans ROB, Bakker SJJ, Kleefstra N, Bilo HJG.** Serum peroxiredoxin 4: a marker of oxidative stress associated with mortality in type 2 diabetes (ZODIAC-28). *PLoS One* 9: e89719, 2014.
14. **Gomez-Cabrera MC, Domenech E, Romagnoli M, Arduini A, Borrás C, Pallardo F V., Sastre J, Viña J.** Oral administration of vitamin C decreases muscle mitochondrial biogenesis and hampers training-induced adaptations in endurance performance. [Online]. *Am J Clin Nutr* 87: 142–9, 2008.  
<http://www.ncbi.nlm.nih.gov/pubmed/18175748>.
15. **Gottfredsen, R.H., Goldstrohm, D.A., Hartney, J.M., Larsen, U.G., Bowler, R.P., Petersen SV.** The cellular distribution of extracellular superoxide dismutase in macrophages is altered by cellular activation but unaffected by the naturally occurring R213G substitution. *Free Radic Biol Med* 69: 348–356, 2014.
16. **Gottfredsen RH, Tran SM, Larsen UG, Madsen P, Nielsen MS, Enghild JJ, Petersen SV.** The C-terminal proteolytic processing of extracellular superoxide dismutase is redox regulated. *Free Radic Biol Med* 52: 191–197, 2012.
17. **Groussard C, Rannou-Bekono F, Machefer G, Chevanne M, Vincent S, Sergent O, Cillard J, Gratas-Delamarche a.** Changes in blood lipid peroxidation markers and antioxidants after a single sprint anaerobic exercise. *Eur J Appl Physiol* 89: 14–20, 2003.
18. **Hara T, Kondo N, Nakamura H, Okuyama H, Mitsui A, Hoshino Y, Yodoi J.** Cell-surface thioredoxin-1: possible involvement in thiol-mediated leukocyte-endothelial cell interaction through lipid rafts. *Antioxid Redox Signal* 9: 1427–37, 2007.

19. **Hitomi Y, Watanabe S, Kizaki T, Sakurai T, Takemasa T, Haga S, Ookawara T, Suzuki K, Ohno H.** Acute exercise increases expression of extracellular superoxide dismutase in skeletal muscle and the aorta. *Redox Rep* 13: 213–216, 2008.
20. **Iversen MB, Gottfredsen RH, Larsen UG, Enghild JJ, Praetorius J, Borregaard N, Petersen S V.** Extracellular superoxide dismutase is present in secretory vesicles of human neutrophils and released upon stimulation. *Free Radic Biol Med* 97: 478–488, 2016.
21. **Jin D-Y, Chae HZ, Rhee SG, Jeang K-T.** Regulatory Role for a Novel Human Thioredoxin Peroxidase in NF- B Activation. *J Biol Chem* 272: 30952–30961, 1997.
22. **Knoops B, Argyropoulou V, Becker S, Ferté L, Kuznetsova O.** Multiple Roles of Peroxiredoxins in Inflammation. *Mol Cells* 39: 60–64, 2016.
23. **Kondo N, Ishii Y, Kwon YW, Tanito M, Sakakura-Nishiyama J, Mochizuki M, Maeda M, Suzuki S, Kojima M, Kim YC, Son A, Nakamura H, Yodoi J. Kondo N1, Ishii Y, Kwon YW, Tanito M, Sakakura-Nishiyama J, Mochizuki M, Maeda M, Suzuki S, Kojima M, Kim YC, Son A, N YJ.** Lipid raft-mediated uptake of cysteine-modified thioredoxin-1: apoptosis enhancement by inhibiting the endogenous thioredoxin-1. *Antioxid Redox Signal* 9: 1439–48, 2007.
24. **Kunze A, Zierath D, Tanzi P, Cain K, Becker K.** Peroxiredoxin 5 (PRX5) Is Correlated Inversely to Systemic Markers of Inflammation in Acute Stroke. *Stroke* 45: 608–10, 2014.
25. **Lamb GD, Westerblad H.** Acute effects of reactive oxygen and nitrogen species on the contractile function of skeletal muscle. *J Physiol* 589: 2119–2127, 2011.
26. **Lee KP, Shin YJ, Cho SC, Lee SM, Bahn YJ, Kim JY, Kwon ES, Jeong do Y, Park S., Rhee SG, Woo HA, Kwon KS.** Peroxiredoxin 3 has a crucial role in the contractile function of skeletal muscle by regulating mitochondrial homeostasis. *Free Radic Biol Med* 77: 298–306, 2014.
27. **Liu CH, Kuo SW, Hsu LM, Huang SC, Wang CH, Tsai PR.** Peroxiredoxin 1 induces inflammatory cytokine response and predicts outcome of cardiogenic shock patients necessitating extracorporeal membrane oxygenation : an observational cohort study and translational approach. *J. Transl. Med.* (2016). doi: 10.1186/s12967-016-

0869-x.

28. **Manabe Y, Takagi M, Nakamura-Yamada M, Goto-Inoue N, Taoka M, Isobe T, Fujii NL.** Redox proteins are constitutively secreted by skeletal muscle. *J Physiol Sci* 64: 401–409, 2014.
29. **Marumoto M, Suzuki S, Hosono A, Arakawa K, Shibata K, Fuku M, Goto C, Tokudome Y, Hoshino H, Imaeda N, Kobayashi M, Yodoi J, Tokudome S.** Changes in thioredoxin concentrations: an observation in an ultra-marathon race. *Env Heal Prev Med* 15: 129–134, 2010.
30. **McArdle F, Pattwell DM, Vasilaki A, McArdle A, Jackson MJ.** Intracellular generation of reactive oxygen species by contracting skeletal muscle cells. *Free Radic Biol Med* 39: 651–7, 2005.
31. **Michailidis Y, Jamurtas AZ, Nikolaidis MG, Fatouros IG, Koutedakis Y, Papassotiriou I, Kouretas D.** Sampling time is crucial for measurement of aerobic exercise-induced oxidative stress. *Med Sci Sport Exerc* 39: 1107–1113, 2007.
32. **Nameta M, Saijo Y, Ohmoto Y, Katsuragi K, Yamamoto K, Yamamoto T, Ishibashi K, Sasaki S.** Disruption of membranes of extracellular vesicles is necessary for ELISA determination of urine AQP2: Proof of disruption and epitopes of AQP2 antibodies. *Int J Mol Sci* 17: 1–7, 2016.
33. **Pattwell D, Ashton T, McArdle A, Griffiths RD, Jackson MJ.** Ischemia and reperfusion of skeletal muscle lead to the appearance of a stable lipid free radical in the circulation. *Am J Physiol Hear Circ Physiol* 284: H2400-4, 2003.
34. **Pattwell DM, McArdle A, Morgan JE, Patridge TA, Jackson MJ.** Release of reactive oxygen and nitrogen species from contracting skeletal muscle cells. *Free Radic Biol Med* 37: 1064–1072, 2004.
35. **Powers SK, Nelson WB, Hudson MB.** Exercise-induced oxidative stress in humans: Cause and consequences. *Free Radic Biol Med* 51: 942–950, 2011.
36. **Powers SK, Talbert EE, Adhietty PJ.** Reactive oxygen and nitrogen species as intracellular signals in skeletal muscle. *J Physiol* 589: 2129–2138, 2011.
37. **Poynton RA, Hampton MB.** Peroxiredoxins as biomarkers of oxidative stress. *Biochim Biophys Acta* 1840: 906–12, 2014.

38. **Riddell JR, Wang XY, Minderman H, Gollnick SO.** Peroxiredoxin 1 stimulates secretion of proinflammatory cytokines by binding to TLR4. *J Immunol* 184: 1022–30, 2010.
39. **Ristow M, Birringer M, Kiehntopf M, Zarse K, Oberbach A, Klo N, Stumvoll M, Kahn CR, Blu M.** Antioxidants prevent health-promoting effects. *PNAS* 106: 8865–8670, 2009.
40. **Salzano S, Checconi P, Hanschmann E-M, Lillig CH, Bowler LD, Chan P, Vaudry D, Mengozzi M, Coppo L, Sacre S, Atkuri KR, Sahaf B, Herzenberg L a, Herzenberg L a, Mullen L, Ghezzi P.** Linkage of inflammation and oxidative stress via release of glutathionylated peroxiredoxin-2, which acts as a danger signal. *Proc Natl Acad Sci U S A* 111: 12157–62, 2014.
41. **Shing CM, Peake JM, Ahern SM, Strobel NA, Wilson G, Jenkins DG, Coombes JS.** The effect of consecutive days of exercise on markers of oxidative stress. *Appl Physiol Nutr Metab* 32, 2007.
42. **Strålin P, Karlsson K, Johansson BO, Marklund SL.** The Interstitium of the Human Arterial Wall Contains Very Large Amounts of Extracellular Superoxide Dismutase. *Arterioscler Thromb Vasc Biol* 15: 2032–2036, 1995.
43. **Szabó-Taylor KÉ, Eggleton P, Turner CAL, Faro ML Lo, Tarr JM, Tóth S, Whiteman M, Haigh RC, Littlechild J a, Winyard PG.** Lymphocytes from rheumatoid arthritis patients have elevated levels of intracellular peroxiredoxin 2, and a greater frequency of cells with exofacial peroxiredoxin 2, compared with healthy human lymphocytes. [Online]. *Int J Biochem Cell Biol* 44: 1223–31, 2012. <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3425769&tool=pmcentrez&rendertype=abstract> [19 Sep. 2013].
44. **They C, Boussac M, Veron P, Ricciardi-Castagnoli P, Raposo G, Garin J, Amigorena S.** Proteomic Analysis of Dendritic Cell-Derived Exosomes: A Secreted Subcellular Compartment Distinct from Apoptotic Vesicles. *J Immunol* 166: 7309–7318, 2001.
45. **Thierry Lévillard and Najate Aït-Ali.** Cell Signaling with Extracellular Thioredoxin and Thioredoxin- Like Proteins: Insight into Their Mechanisms of Action. *Oxid Med Cell Longev* 2017: 11 pages, 2017.

46. **Turner JE, Bennett SJ, Campbell JP, Bosch JA, Aldred S, Griffiths HR.** The antioxidant enzyme peroxiredoxin-2 is depleted in lymphocytes seven days after ultra-endurance exercise. [Online]. *Free Radic Res* 47: 821–8, 2013.  
<http://www.ncbi.nlm.nih.gov/pubmed/23889121> [19 Sep. 2013].
47. **Wadley AJ, Chen YW, Bennett SJ, Lip GYH, Turner JE, Fisher J., Aldred S.** Monitoring changes in thioredoxin and over-oxidised peroxiredoxin in response to exercise in humans. *Free Radic Res* 49: 290–298, 2015.
48. **Wadley AJ, Chen YW, Lip GY., Fisher JP, Aldred S.** Low volume-high intensity interval exercise elicits anti-oxidant and anti-inflammatory effects in humans. *J Sports Sci* 27: 1–9, 2015.
49. **Wadley AJ, Holliday A, Morgan RG, Heesom KJ, Aldred S, Peters DM, Bueno AA, Coles SJ.** Preliminary evidence of reductive stress in human cytotoxic T-cells following exercise. *J. Appl. Physiol.* .
50. **Winterbourn CC.** The biological chemistry of hydrogen peroxide. 1st ed. Elsevier Inc.

## Tables

**Table 1.** Demographics for participants in studies 1 and 2.

	Energy-matched Trials (study 1)	Eccentric-based Resistance Exercise (study 2)	Statistical Analysis
<b>Number of Participants</b>	9	16	n/a
<b>Age (years)</b>	29 ±	25 ±	P = NS
	5	9	
<b>Body Mass Index (kg/m<sup>2</sup>)</b>	24.2 ±	25.3 ±	P = NS
	3.4	4.1	
<b>IPAQ (METs- min/week)</b>	6683 ±	2540 ±	+P = 0.005
	3835*	2022	+P = 0.004
<b>Watt Max (Watt/kg)</b>	3.4 ±		n/a
	0.5		
<b><math>\dot{V}O_2</math><sub>MAX</sub> (mL.kg<sup>-1</sup>.min<sup>-1</sup>)</b>	44.5 ±		n/a
	6.4		

Grey boxes indicate missing data.

\* Indicates significant difference in comparison to study 2: \*P < 0.05, \*\*P < 0.001.

NS P > 0.05.

**Table 2.** Physiological response to aerobic-based exercise (study 1).

Energy-matched Cycling Trials (Study 1)			Statistical Analysis
Trial	Continuous cycling for 58 min, predicted	10 x 4 min cycling intervals, predicted	--
	60% $\dot{V}O_{2\text{ MAX}}$ (MOD)	85% $\dot{V}O_{2\text{ MAX}}$ (2 min rest intervals. Total time = 58 min, HIIE)	
Mean $\dot{V}O_{2\text{ MAX}}$ (%)	56.5 ± 2.6	58.9 ± 4.3	P = NS
Energy Expenditure (kJ)	2077 ± 340	2072 ± 339	P = NS
Average RPE	12 ± 1	16 ± 1 <sup>***</sup>	<sup>***</sup> P < 0.0001
Mean Blood Lactate (mmol/L)	1.9 ± 0.6	6.8 ± 1.4	<sup>***</sup> P < 0.0001
Mean Blood Glucose (mmol/L)	3.9 ± 0.3	4.5 ± 0.6	P = NS

\* Indicates a significant difference between MOD and HIIE: <sup>\*\*\*</sup> P < 0.0001.

NS P > 0.05.

Table 3. Changes in markers or muscle damage following eccentric-based resistance exercise (study 2). Values are means  $\pm$  standard deviation.

	Pre	Post+0	Post+30min	Post+3hr	Post+48hr
Creatine Kinase	147.6 $\pm$	236.1 $\pm$	289.9 $\pm$	560.8 $\pm$	575.9 $\pm$
(Units/L)	27.1	65.5 *	86.0 *+	273.5 **+#	290.8 **+#\$
Lactate	254.9 $\pm$	282.7 $\pm$	274.1 $\pm$	290.3 $\pm$	299.9 $\pm$
Dehydrogenase	130.6	70.9 *	77.1 *	77.8 *+	165.2 *+
(Units/L)					

\* Indicates significant difference in comparison to Pre: \*P < 0.05, \*\*P < 0.001.

+ Indicates significant difference in comparison to Post+0: \*P < 0.05.

# Indicates significant difference in comparison to Post+30min: #P < 0.05.

\$ Indicates significant difference in comparison to Post+3hr: \$P < 0.05.

## Figures

**Figure 1:** Schematic of the two exercise studies. Dark lines represent the exercise session, with lighter lines indicating pre- and post-exercise resting periods. Gaps between dark lines indicate the rest periods during the HIIE trial. Blood samples taken for each study are indicated as arrows.

**Figure 2:** Changes in redox enzyme concentration in response to two energy-matched cycling bouts (A) - moderate steady state (MOD - black bars) and high intensity interval exercise (HIIE - white bars) and an eccentric-based resistance exercise protocol (B): PRDX-2, PRDX-4, TRX-1, TRX-R and SOD3. Values are means  $\pm$  standard error. For Figure 2A: \* indicates significant differences relative to Pre: \*  $p < .05$ . # indicates a significant difference relative to Post+0: #  $p < .05$ . \$ indicates a significant difference relative to Post+30: \$  $p < .05$ . + indicates a significant difference between MOD and HIIE: +  $p < .05$ . For Figure 2B: ^ indicates a significant difference between Post+30min and Post+48hrs timepoints.

Figure 3: Changes in plasma IL-6 in response to two energy-matched cycling bouts (A): moderate steady state (MOD - black bars) and high interval exercise (HIIE - white bars) and an eccentric-based resistance exercise protocol (B). Values are means  $\pm$  standard error. For Figures 3A and 3B: \* indicates significant differences relative to Pre: \*  $p < .05$ ; \*\*  $p < .001$ . # indicates a significant difference relative to Post+0: #  $p < .05$ . \$ indicates a significant difference relative to Post+30: \$  $p < .05$ . + indicates a significant difference between MOD and HIIE: +  $p < .05$ .