



## Genome editing and regeneration pipeline for engineering disease resistance in tomato using CRISPR/Cas9

Item Type	Article (Version of Record)
UoW Affiliated Authors	Tör, M.
Full Citation	Erdoğan, İ., Debbarma, R., Sherry, M., Mancak, İ., Grant, M. and Tör, M. (2026) Genome editing and regeneration pipeline for engineering disease resistance in tomato using CRISPR/Cas9. <i>Frontiers in Plant Science</i> , 17. pp. 1-10. ISSN 1664-462X
DOI/ISBN/ISSN	<a href="https://doi.org/10.3389/fpls.2026.1754287">https://doi.org/10.3389/fpls.2026.1754287</a>
Journal/Publisher	Frontiers in Plant Science Frontiers
Rights/Publisher Set Statement	© 2026 Erdoğan, Debbarma, Sherry, Mancak, Grant and Tör. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY), <a href="https://creativecommons.org/licenses/by/4.0/">https://creativecommons.org/licenses/by/4.0/</a>
License	CC BY 4.0
Link	<a href="https://www.frontiersin.org/journals/plant-science/articles/10.3389/fpls.2026.1754287/full">https://www.frontiersin.org/journals/plant-science/articles/10.3389/fpls.2026.1754287/full</a>

For more information, please contact [wrapteam@worc.ac.uk](mailto:wrapteam@worc.ac.uk)



## OPEN ACCESS

### EDITED BY

Ahmad A. Omar,  
University of Florida, United States

### REVIEWED BY

Rashmi Kaul,  
International Centre for Genetic  
Engineering and Biotechnology, India  
Fernanda dos Santos Nascimento,  
Brazilian Agricultural Research  
Corporation (EMBRAPA), Brazil  
Johni Debbarma,  
Assam Don Bosco University, India

### \*CORRESPONDENCE

Mahmut Tör  
✉ m.tor@worc.ac.uk

<sup>†</sup>These authors have contributed equally  
to this work

RECEIVED 25 November 2025

REVISED 18 February 2026

ACCEPTED 23 February 2026

PUBLISHED 24 March 2026

### CITATION

Erdoğan İ, Debbarma R, Sherry M,  
Mancak İ, Grant M and Tör M (2026)  
Genome editing and regeneration  
pipeline for engineering disease  
resistance in tomato using CRISPR/Cas9.  
*Front. Plant Sci.* 17:1754287.  
doi: 10.3389/fpls.2026.1754287

### COPYRIGHT

© 2026 Erdoğan, Debbarma, Sherry,  
Mancak, Grant and Tör. This is an open-  
access article distributed under the terms  
of the [Creative Commons Attribution  
License \(CC BY\)](https://creativecommons.org/licenses/by/4.0/). The use, distribution or  
reproduction in other forums is  
permitted, provided the original  
author(s) and the copyright owner(s) are  
credited and that the original publication  
in this journal is cited, in accordance  
with accepted academic practice. No  
use, distribution or reproduction is  
permitted which does not comply with  
these terms.

# Genome editing and regeneration pipeline for engineering disease resistance in tomato using CRISPR/Cas9

İbrahim Erdoğan <sup>1†</sup>, Ruma Debbarma <sup>1†</sup>, Minu Sherry <sup>1</sup>,  
İbrahim Mancak <sup>2</sup>, Murray Grant <sup>1</sup> and Mahmut Tör <sup>3\*</sup>

<sup>1</sup>University of Warwick, School of Life Sciences, Elizabeth Creak Horticultural Technology Centre (ECHTC), Coventry, United Kingdom, <sup>2</sup>Manier Seeds, Adana, Türkiye, <sup>3</sup>Molecular Plant and Microbial Biosciences Research Unit, School of Science and the Environment, University of Worcester, Worcester, United Kingdom

CRISPR/Cas9 (Clustered Regularly Interspaced Short Palindromic Repeats/Cas9)-based genome editing has emerged as a powerful tool for developing disease-resistant crops. Here, we present a comprehensive and reproducible protocol for applying CRISPR/Cas9 genome editing in tomato (*Solanum lycopersicum*), covering guide RNA (gRNA) design using CRISPOR, Golden Gate vector assembly, *Agrobacterium*-mediated transformation, plant regeneration, and molecular validation of edited plants. The workflow integrates standardized bioinformatics and sequencing-based validation tools, including DSDecodeM, TIDE, and protein-level impact analysis, to confirm targeted mutations and assess editing efficiency. Quantitative benchmarks for regeneration, transformation, and editing efficiencies were provided to support reproducibility. This protocol offers an integrated pipeline for generating and validating targeted gene knockouts in tomatoes and is intended to facilitate functional genomic studies and the development of disease-resistant cultivars. However, it is more widely applicable to gene editing in tomato plants.

### KEYWORDS

crop improvement, disease resistance, genome editing, susceptibility gene, targeted mutations

## 1 Introduction

Tomato (*Solanum lycopersicum*) is one of the most important vegetable crops worldwide, contributing substantially to global vegetable production and human nutrition. It is a major dietary source of carotenoids, particularly lycopene and  $\beta$ -carotene, which are associated with important health benefits. However, tomato production is frequently limited by a wide range of bacterial, oomycete, fungal, and viral pathogens that cause significant yield loss and economic damage. Global tomato production continues to expand to meet the demand from both fresh markets and processing industries (WPTC; Kaul et al., 2025), increasing the need for durable and sustainable disease resistance strategies.

Conventional breeding approaches for disease resistance are often slow and constrained by limited genetic diversity and linkage drags. CRISPR/Cas9 (Clustered Regularly Interspaced Short Palindromic Repeats/Cas9) genome editing is a precise and efficient

alternative for introducing targeted genetic modifications (Erdoğan et al., 2023; Tiwari et al., 2023). This technology enables the rapid functional analysis of genes and the development of improved cultivars with desirable agronomic traits. CRISPR/Cas9 has been widely applied in tomatoes to modify traits, including disease resistance, demonstrating its versatility as a tool for crop improvement (Thomazella et al., 2021; Nekrasov et al., 2017; Tashkandi et al., 2018; Ortigosa et al., 2019; Ortega et al., 2024; Sandhya et al., 2025).

Here, we describe a detailed and reproducible protocol for generating disease-resistant tomato plants through targeted editing of *susceptibility* (*S*) genes using the CRISPR/Cas9 system. The protocol integrates guide RNA (gRNA) design and off-target prediction using CRISPOR, Golden Gate cloning, *Agrobacterium*-mediated transformation, optimized tissue culture-based regeneration, and standardized molecular validation of edited plants. By combining optimized regeneration procedures with quantitative benchmarks and sequencing-based mutation analysis, this workflow provides an integrated and experimentally validated pipeline for producing targeted gene knockouts. This protocol is designed to support both functional genomics research and the practical implementation of genome-editing strategies for developing improved tomato varieties.

## 2 Materials and methods

### 2.1 Materials

#### 2.1.1 Laboratory supplies and consumables

Pipette sets and sterile tips, 0.5- and 1.5-mL microcentrifuge tubes, PCR tubes, disposable plate spreaders and inoculation loops, tube racks, sterile distilled water, ice buckets, personal protective equipment (gloves and masks), and standard laboratory disinfectants were used throughout the experiments.

#### 2.1.2 Reagents and media

LB broth and LB agar, antibiotics (including spectinomycin, ampicillin, chloramphenicol, kanamycin, rifampicin, and gentamicin), Murashige and Skoog (MS) medium, plant growth regulators (IAA, BAP, zeatin, and IBA), timentin, agar (0.7%), SOC medium, and acetosyringone (ACS) were used.

**Abbreviations:** CRISPR, Clustered Regularly Interspaced Short Palindromic Repeats; Cas9, CRISPR-associated protein 9; gRNA, Guide RNA; sgRNA, Single-guide RNA; MS, Murashige and Skoog medium; IAA, Indole-3-acetic acid; LB, Luria-Bertani medium; PCR, Polymerase Chain Reaction; DNA, Deoxyribonucleic Acid; RNA, Ribonucleic Acid; bp, Base pair; kb, Kilobase; OD, Optical Density; NPTII, Neomycin Phosphotransferase II; dNTPs, Deoxynucleotide Triphosphates; NFW, Nuclease-Free Water; HRM, High-Resolution Melting; T0/T1, Transgenic Generation 0/1; WT, Wild Type; A. tumefaciens, *Agrobacterium tumefaciens*; E. coli, *Escherichia coli*; PGRs, Plant Growth Regulators.

#### 2.1.3 Molecular biology reagents

BpiI and BsaI restriction enzymes, T4 DNA ligase, plasmid DNA isolation kits, PCR reagents and primers, and Sanger DNA sequencing services were used for molecular cloning and validation of the constructs.

#### 2.1.4 Biological materials

Tomato (*S. lycopersicum*) cv. “Ailsa Craig” seeds, CRISPR/Cas9 vectors (pDGE5, pDGE8, and pDGE1), *Agrobacterium tumefaciens* strain GV3101, electrocompetent *Escherichia coli* DH10b cells, and electroporation cuvettes were used.

#### 2.1.5 Equipment

A thermal cycler (PCR machine), BioRad MicroPulser electroporator, centrifuge, and controlled-environment growth chambers were used.

### 2.2 gRNA design and vector construction

gRNAs targeting the tomato susceptibility gene *MLO1* were identified using the NCBI database (O’Leary et al., 2024). gRNAs were designed using the CRISPOR web tool (Concordet and Haeussler, 2018) based on the presence of PAM, GC content, and predicted specificity scores. Potential off-target sites were evaluated *in silico* using the tomato reference genome. The overall workflow used for CRISPR/Cas9-mediated genome editing and plant regeneration is illustrated in Figure 1.

gRNA sequences:

gRNA1 Forward: ATTGGGAGGTACCACGCAATGGTG.

gRNA1 Reverse: AAACCACCATTGCGTGGTACCTCC.

gRNA2 Forward: ATTGCCATGGTTAGCCTTATGGCT.

gRNA2 Reverse: AAACAGCCATAAGGCTAACCATGG.

#### 2.2.1 Oligonucleotide hybridization

Lyophilized oligonucleotides were resuspended to 100  $\mu$ M in sterile water and then diluted to 10  $\mu$ M. Equal volumes (5  $\mu$ L each) of forward and reverse oligonucleotides were mixed with 40  $\mu$ L of sterile water to obtain a 5  $\mu$ M hybridization solution. The mixture was denatured at 98  $^{\circ}$ C for 5 min and allowed to cool to room temperature (RT). A 1:100 dilution (50 fmol/ $\mu$ L) was prepared for cloning and stored at  $-20^{\circ}$ C.

#### 2.2.2 Golden Gate cloning

gRNA1 and gRNA2 were first cloned into shuttle vectors pDGE5 and pDGE8, respectively, and subsequently assembled into the final binary vector pDGE1 using Golden Gate cloning (Ordon et al., 2017; Stuttmann et al., 2021). The Golden Gate cloning strategy used for gRNA assembly into the transformation vector is illustrated in Figure 2, and the plasmid constructs used for genome editing are shown in Figure 3.

Cut–ligation reaction for shuttle vectors (10  $\mu$ L total volume):

- 4  $\mu$ L of dH<sub>2</sub>O.
- 3  $\mu$ L of pDGE5 or pDGE8 (20 fmol  $\approx$  200 ng).
- 1  $\mu$ L of hybridized oligonucleotides.
- 1  $\mu$ L of 10 $\times$  ligation buffer.
- 0.5  $\mu$ L of BpiI (10U/ $\mu$ L).
- 0.5  $\mu$ L of T4 DNA ligase (10U/ $\mu$ L).

Final vector assembly (20  $\mu$ L total volume):

- 11.5  $\mu$ L of dH<sub>2</sub>O.
- 3.5  $\mu$ L of pDGE1 (200 ng).
- 0.5  $\mu$ L of pDGE5-gRNA1 (20 fmol  $\approx$  220 ng).
- 0.5  $\mu$ L of pDGE8-gRNA2 (20 fmol  $\approx$  220 ng).
- 2  $\mu$ L of 10 $\times$  ligation buffer.
- 1  $\mu$ L of BsaI (10U/ $\mu$ L).
- 1  $\mu$ L of T4 DNA ligase (10U/ $\mu$ L).

The thermocycling conditions were as follows: 37  $^{\circ}$ C for 2 min and 16  $^{\circ}$ C for 5 min (10–30 cycles), followed by 50  $^{\circ}$ C for 10 min and 80  $^{\circ}$ C for 10 min. The reaction products were purified using spin columns and stored at 4  $^{\circ}$ C.

### 2.2.3 Transformation into *E. coli*

Electrocompetent *E. coli* DH10b cells (15  $\mu$ L) were mixed with 3  $\mu$ L of purified ligation product and electroporated at 2.5 kV (cuvette gap: 0.1 cm). Cells were recovered in 250  $\mu$ L of SOC medium at 37 $^{\circ}$ C for 1 h with shaking and plated on LB agar containing spectinomycin. Single colonies were cultured overnight, and plasmid DNA was isolated using a miniprep kit.

## 2.3 *Agrobacterium*-mediated transformation

Purified plasmids were introduced into electrocompetent *A. tumefaciens* GV3101 by electroporation. Cells were recovered in SOC medium at 28  $^{\circ}$ C for 1 h and plated on LB agar containing spectinomycin, gentamicin, and rifampicin. The colonies were cultured in selective LB broth and stored as glycerol stocks.

## 2.4 Tomato regeneration and selection

The tomato regeneration and transformation pipeline used in this study is summarized in Figure 4. Sterilized tomato seeds were germinated *in vitro* on full-strength MS medium with 0.7% agar for 10 days. Cotyledon explants were co-cultivated with *Agrobacterium* suspension (OD<sub>600</sub> 0.3–0.5) supplemented with 100  $\mu$ M ACS for 5–

10 min and incubated on co-cultivation medium in the dark for 48 h.

The explants were transferred to shoot induction medium (TS1), which consisted of full-strength MS supplemented with zeatin (0.5 mg/L), IAA (0.1 mg/L), BAP (0.1 mg/L), kanamycin (10 mg/L), and timentin (200 mg/L). The cultures were maintained under a 16-h light/8-h dark photoperiod. Shoots emerged approximately 2 weeks after co-cultivation and were maintained on TS1 medium for approximately 7 weeks.

Regenerated shoots were transferred to half-strength MS rooting medium supplemented with IBA and subsequently acclimatized in soil under controlled growth chamber conditions (23  $^{\circ}$ C, long-day photoperiod) (Figure 5).

Regeneration, transformation, and editing efficiencies were calculated based on the number of co-cultivated explants and confirmed edited plants.

## 2.5 gRNA off-target analysis

Off-target predictions were performed using CRISPOR. Candidate sites with  $\leq 3$  mismatches and canonical PAM sequences were evaluated. The specificity scores and predicted off-target counts are summarized in Table 1.

## 2.6 Molecular screening and genome editing validation

Genomic DNA was extracted from the regenerated plants using a modified Shorty DNA extraction protocol (Edwards et al., 1991). DNA quality was assessed using agarose gel electrophoresis.

Genome editing in kanamycin-resistant T0 plants was validated by PCR amplification of the target locus, followed by Sanger sequencing. Sequence traces were analyzed using DSDecodeM and TIDE to determine the indel patterns and mutation frequencies. Predicted protein consequences were evaluated using the ExPASy Translate tool, which revealed frameshift mutations and premature stop codons in the edited lines.


# 3 Results

## 3.1 PCR confirmation of CRISPR/Cas9 integration

The integration of the CRISPR/Cas9 construct into the regenerated tomato plants was verified using PCR with kanamycin-specific primers [Forward: CTTCCCCTTCAGTGACAAC, Reverse: TTGGGTGGAGAGGCTATTCC]. Amplification of the expected fragment was observed in kanamycin-positive transformants (kan<sup>+</sup> PT), whereas wild-type (WT) plants showed

TABLE 1 Predicted off-target specificity metrics and off-target counts for guide RNAs targeting MLO1 locus.

Position/Strand	Guide sequence + PAM	mitSpecScore	cfdSpecScore	Off-targets count
329rev	GGAGGTACCACGCAATGGTGTGG	100	100	0
111rev	CCATGGTTAGCCTTATGGCTAGG	95	98	12



**FIGURE 1**  
CRISPR/Cas9 workflow in tomato. Schematic overview of the integrated genome editing workflow, including gRNA design using CRISPOR, Golden Gate vector assembly, *Agrobacterium*-mediated transformation, plant regeneration, and molecular validation of edited plants.

no band (Figure 6). These results confirmed the successful T-DNA integration in the selected T<sub>0</sub> plants.


PCR amplification was performed in 25- $\mu$ L reaction volumes. Each reaction contained 12.5  $\mu$ L of Red Mix, 0.5  $\mu$ L of forward primer, 0.5  $\mu$ L of reverse primer, and nuclease-free dH<sub>2</sub>O adjusted to volume, with 1  $\mu$ L of template DNA added per reaction. Reactions were prepared as a master mix for 10 samples to minimize the pipetting variability. PCR was carried out under the following cycling conditions: initial denaturation at 95°C for 1 min;

30 cycles of denaturation at 95°C for 15 s, annealing at 60°C for 15 s, and extension at 72°C for 10s; followed by a final extension at 72°C for 5 min. Amplified products were stored at 4°C until further analysis.


### 3.2 Sequence-based detection of indels

To detect the targeted mutations, genomic regions flanking the gRNA target sites were amplified using the following primers:

### Golden Gate Cloning gRNA Assembly Workflow



**FIGURE 2**  
Golden Gate cloning strategy. Diagram illustrating the assembly of gRNA constructs using pDGE5 and pDGE8 shuttle vectors, followed by integration into the final pDGE1 binary vector (Ordon et al., 2017; Stuttmann et al., 2021).



Forward: ATGATCAGTGGAGGCATGCT.  
Reverse: GGAGTTGGGTAAGGAGTTGGA.

Targeted regions flanking the gRNA sequences were PCR-amplified and Sanger-sequenced. Analysis with DSDecodeM revealed various indel patterns in the kanamycin-positive plants. Representative alleles of *MLO1* are shown in Figure 7.

### 3.3 TIDE analysis of indel frequency

Indel frequencies were further quantified using TIDE, which decomposes sequencing traces to determine insertion/deletion distributions and mutation efficiencies. TIDE analysis (Figure 8) indicated editing efficiencies ranging from 40% to 70% in T<sub>0</sub> plants, consistent with PCR and DSDecodeM data. These data demonstrate the reproducibility of the workflow across multiple transformants.

### 3.4 Predicted protein consequences

Translation of the edited alleles using the ExpASY Translate tool revealed frameshift mutations that introduced premature stop codons (Figure 9). These mutations were predicted to truncate the encoded proteins, consistent with the generation of loss-of-function alleles in *MLO1*.

### 3.5 Transformation, regeneration, and editing efficiency

Table 2 summarizes the number of co-cultivated explants, regeneration, transformation efficiency based on Kan diagnostic PCR, and editing efficiency of the transformants.

Table 1 lists the gRNA specificity scores and predicted off-target counts from the CRISPOR analysis. No high-probability off-target sites were detected for gRNA1, whereas gRNA2 had minimal

## TOMATO PLANT REGENERATION PIPELINE




FIGURE 4

Tomato regeneration pipeline. Sequential stages of tomato transformation and regeneration following *Agrobacterium*-mediated gene transfer. (A) *In vitro* seed germination. (B) Co-cultivation of cotyledon explants with *Agrobacterium*. (C) Shoot regeneration on TS1 medium. (D) Shoot development. (E) Root induction. (F) Acclimatization and hardening of regenerated plants in soil.





FIGURE 5

Stage-wise regeneration of tomato plants. (A) *In vitro* seed germination; (B) co-cultivation of cotyledon explants; (C) callus formation and shoot regeneration; (D) shoot development on TS1 medium; (E) root induction; (F) acclimatization of regenerated plantlets.



**FIGURE 6**  
 PCR confirmation of putative transformants. Examples of PCR amplification using kanamycin-specific primers to confirm construct integration. M, DNA marker; kan<sup>+</sup> PT, kanamycin-positive putative transformants; WT, wild-type Ailsa Craig.

predicted off-targets (≤12 with low specificity risk), supporting the high precision of the designed gRNAs, ranked by default from highest to lowest specificity score (Hsu et al., 2013).

## 4 Discussion

### 4.1 Methodological advancement

This protocol comprises a facile process for groups to establish gene-editing experiments in tomato plants. It integrates optimized regeneration conditions, Golden Gate cloning, molecular validation (PCR and sequencing), and bioinformatic assessment (DSDecodeM, TIDE, and ExPASy) into a reproducible pipeline for CRISPR/Cas9-mediated genome editing in tomato. Quantitative benchmarks for transformation, regeneration, and editing efficiency allow researchers to replicate and adapt this workflow.

### 4.2 Reproducibility and validation


The combination of PCR confirmation, sequencing-based indel detection, and TIDE analysis provides the user with a robust, multilayered validation strategy. Predicted protein-level consequences ensure that the observed mutations are functionally relevant, enabling reliable knockout generation.

### 4.3 Expected outcomes

- Shoot regeneration occurs within 2–3 weeks post-transformation, and rooted plantlets are ready for acclimatization in 6–8 weeks.
- Editing efficiencies typically range from 40% to 70%, depending on the gRNA design and tomato genotype.
- Homozygous or biallelic T<sub>0</sub> plants displayed frameshift mutations and premature stop codons, consistent with loss of function.
- Transgene-free edited lines can be obtained in subsequent generations by segregation.
- Phenotypic assays under pathogen challenge (powdery mildew, *Oidium neolycopersici*) are expected to demonstrate enhanced disease resistance in knockout lines compared to that in the WT.

### 4.4 Best practices

- Select gRNAs with high predicted efficiency and minimal off-target effects.
- Maintain strict aseptic technique during tissue culture.
- Always include WT and empty vector controls at all stages of the experiment.
- Whole-genome sequencing may be considered for comprehensive off-target analyses if end objectives are required.



**FIGURE 7**  
 Mutation analysis using DSDecodeM. Representative deletion patterns in kanamycin-positive plants. AC-WT: wild-type sequence; A1 and A2: edited alleles. Blue sequences indicate the gRNA target sites. Deletions are represented by dashed lines. Biallelic mutations were observed in several lines, indicating effective genome editing at the target locus.





TABLE 2 Regeneration, transformation, and editing efficiencies of the *MLO1* gene-editing construct.

Gene construct	No. of co-cultivated explants	Regeneration efficiency (%)	Transformation efficiency (%)	Editing efficiency (%)
<i>MLO1</i>	75	62.70	37.33	50

### 4.5 Applicability

This workflow is broadly adaptable to other tomato cultivars or related Solanaceae crops with minor modifications. This protocol facilitates functional genomic studies and accelerates

the generation of disease-resistant tomato varieties (Table 3). #Common technical issues encountered during the tomato transformation and regeneration workflow, along with suggested troubleshooting strategies, are summarized in Table 3.

TABLE 3 Troubleshooting.

Step	Problem	Possible cause	Solution
gRNA cloning	Low ligation efficiency	Incomplete digestion or inefficient ligase	Verify enzyme activity, increase ligation time or temperature cycles
<i>E. coli</i> transformation	No colonies after electroporation	Poor competency or electrical settings	Use freshly prepared competent cells, ensure correct electroporation parameters (or provide)?
<i>Agrobacterium</i> transformation	No growth on selective plates	Antibiotic concentrations too high or plasmid lost	Confirm antibiotic stock concentrations, recheck plasmid integrity (sequence)
Shoot regeneration	Poor shoot induction	Suboptimal PGR concentrations or dark period	Optimize Zeatin/IAA/BA ratios, ensure 48 h dark co-cultivation
High contamination	Bacterial/fungal contamination in culture	Poor aseptic technique or non-sterile seeds	Use 70% ethanol and bleach sterilization for seeds, work in laminar flow cabinet
Low editing efficiency	No mutations detected	gRNA not effective or low Cas9 activity	Redesign gRNA targeting exonic conserved regions, verify expression with RT-PCR
Chimerism in T <sub>0</sub> plants	Mosaic editing due to late Cas9 activity	Common in first-generation CRISPR plants	Screen T <sub>1</sub> generation to identify stable lines, ideally vector free lines
No root formation	Shoot death on rooting medium	Hormone imbalance or bacterial overgrowth	Adjust IBA concentration, use fresh media, check for latent <i>Agrobacterium</i>

## Data availability statement

The original contributions presented in the study are included in the article/supplementary material. Further inquiries can be directed to the corresponding author.

## Author contributions

IE: Writing – original draft, Investigation, Data curation, Writing – review & editing, Methodology. RD: Data curation, Writing – review & editing, Investigation. MS: Writing – review & editing, Methodology, Investigation. IM: Conceptualization, Funding acquisition, Writing – review & editing. MG: Conceptualization, Supervision, Resources, Writing – review & editing. MT: Project administration, Writing – original draft, Resources, Supervision, Conceptualization, Writing – review & editing, Funding acquisition.

## Funding

The author(s) declared that financial support was received for this work and/or its publication. Financial support from Manier Seeds grants R210125 to MT is gratefully acknowledged. MG acknowledges the Elizabeth Creak Charitable Trust for infrastructure and financial support for the establishment of the Elizabeth Creak Horticultural Technology Centre. We acknowledge the University of Worcester for covering the article publishing charges (APC).

## Conflict of interest

IM was employed by Manier Seeds.

The remaining authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

The author MT declared that they were an editorial board member of Frontiers, at the time of submission. This had no impact on the peer review process and the final decision.

The authors declare that this study received funding from Manier Seeds. The funder's involvement was limited to provision of consumables and contribution to writing and editing the manuscript through its employee I.M. The funder was not involved in the study design, collection, analysis, or interpretation of data, or the decision to submit it for publication.

## Generative AI statement

The author(s) declared that generative AI was not used in the creation of this manuscript.

Any alternative text (alt text) provided alongside figures in this article has been generated by Frontiers with the support of artificial intelligence and reasonable efforts have been made to ensure accuracy, including review by the authors wherever possible. If you identify any issues, please contact us.

## Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

## References

- Concordet, J. P., and Haeussler, M. (2018). CRISPOR: intuitive guide selection for CRISPR/Cas9 genome editing experiments and screens. *Nucleic Acids Res.* 46, W242–W245. doi: 10.1093/nar/gky354
- Edwards, K., Johnstone, C., and Thompson, C. (1991). A simple and rapid method for the preparation of plant genomic DNA for PCR analysis. *Nucleic Acids Res.* 19, 1349. doi: 10.1093/nar/19.6.1349
- Erdoğan, İ., Cevher-Keskin, B., Bilir, Ö., Hong, Y., and Tör, M. (2023). Recent developments in CRISPR/Cas9 genome-editing technology related to plant disease resistance and abiotic stress tolerance. *Biology* 12, 1037. doi: 10.3390/biology12071037
- Hsu, P. D., Scott, D. A., Weinstein, J. A., Ran, F. A., Konermann, S., Agarwala, V., et al. (2013). DNA targeting specificity of RNA-guided Cas9 nucleases. *Nat. Biotechnol.* 31, 827–832. doi: 10.1038/nbt.2647
- Kaul, R., Thangaraj, A., Sharda, S., and Kaul, T. (2025). Optimization of tissue culture and Cas9 transgene expression in tomato: A step towards CRISPR/Cas9-based genetic improvement. *Plant Sci.* 352, 112324. doi: 10.1016/j.plantsci.2024.112324
- Nekrasov, V., Wang, C., Win, J., Lanz, C., Weigel, D., and Kamoun, S. (2017). Rapid generation of a transgene-free powdery mildew resistant tomato by genome deletion. *Sci. Rep.* 7, 482. doi: 10.1038/s41598-017-00578-x
- O'Leary, N. A., Cox, E., Holmes, J. B., Anderson, W. R., Falk, R., Hem, V., et al. (2024). Exploring and retrieving sequence and metadata for species across the tree of life with NCBI Datasets. *Sci. Data* 11, 732. doi: 10.1038/s41597-024-03571-y
- Ordon, J., Gantner, J., Kemna, J., Schwalgun, L., Reschke, M., Streubel, J., et al. (2017). Generation of chromosomal deletions in dicotyledonous plants employing a user-friendly genome editing toolkit. *Plant J.* 89, 155–168. doi: 10.1111/tbj.13319
- Ortega, A., Seong, K., Schultink, A., de Toledo Thomazella, D. P., Seo, E., Zhang, E., et al. (2024). CRISPR/Cas9-mediated editing of Bs5 and Bs5L in tomato leads to resistance against *Xanthomonas*. *Plant Biotechnol. J.* 22, 2785–2787. doi: 10.1111/pbi.14404PMC
- Ortigosa, A., Gimenez-Ibanez, S., Leonhardt, N., and Solano, R. (2019). Design of a bacterial speck resistant tomato by CRISPR/Cas9-mediated editing of *SlJAZ2*. *Plant Biotechnol. J.* 17, 665–673. doi: 10.1111/pbi.13006
- Sandhya, D., Jogam, P., Sheri, V., Peddaboina, V., Allini, V. R., and Abbagani, S. (2025). Efficient knockout of *TOM1* gene using CRISPR/Cas9 in Indian tomato cultivar 'Arka Vikas' (Springer Nature Singapore, Vegetos), 1–10. doi: 10.1007/s42535-024-01138-1
- Stuttman, J., Barthel, K., Martin, P., Ordon, J., Erickson, J. L., Herr, R., et al. (2021). Highly efficient multiplex editing: one-shot generation of 8× *Nicotiana benthamiana* and 12× *Arabidopsis* mutants. *Plant J.* 106, 8–22. doi: 10.1111/tbj.15197
- Tashkandi, M., Ali, Z., Aljedaani, F., Shami, A., and Mahfouz, M. M. (2018). Engineering resistance against Tomato yellow leaf curl virus via the CRISPR/Cas9 system in tomato. *Plant Signaling Behav.* 13, e1525996. doi: 10.1080/15592324.2018.1525996
- Thomazella, D. P. D. T., et al. (2021). Loss of function of a DMR6 ortholog in tomato confers broadspectrum disease resistance. *Proceedings of the National Academy of Sciences*. doi: 10.1073/pnas.2026152118
- Tiwari, J. K., Singh, A. K., and Behera, T. K. (2023). CRISPR/Cas genome editing in tomato improvement: Advances and applications. *Front. Plant Sci.* 14. doi: 10.3389/fpls.2023.1121209